

k -chromatic number of graphs on surfaces*

Zdeněk Dvořák[†]

Riste Škrekovski[‡]

Abstract

A well-known result (Heawood [6], Ringel [11], Ringel and Youngs [10]) states that the maximum chromatic number of a graph embedded in a given surface S coincides with the size of the largest clique that can be embedded in S , and that this number can be expressed as a simple formula in the Eulerian genus of S . We study maximum chromatic number of k edge-disjoint graphs embedded in a surface. We improve the previously known upper bounds, and show that in many cases, the new upper bound coincides with the lower bound obtained from embedding disjoint cliques in the surface. In the proof of this result, we derive a variant of Euler's Formula for union of several graphs that might be interesting independently.

1 Introduction and Definitions

We consider simple undirected graphs with no loops and parallel edges. Let $e(G)$ and $n(G)$ denote the number of edges and the number of vertices of a graph G , respectively. When the graph G is clear from the context, we simply use e and n . A *proper coloring* of a graph G by k colors is assignment of colors $1, 2, \dots, k$ to vertices of G such that no two adjacent vertices have the same color. The *chromatic number* $\chi(G)$ of graph G is the minimum k such that G has a proper coloring by k colors.

Let Σ_h denote the orientable surface obtained from the sphere by attaching h handles, and let Π_h be the nonorientable surface obtained from the

*Supported in part by bilateral projects SLO-CZ/04-05-002 and MSMT-07-0405 between Slovenia and Czech Republic.

[†]Charles University, Faculty of Mathematics and Physics, Institute for Theoretical Computer Science (ITI), Malostranské nám. 2/25, 118 00, Prague, Czech Republic, rakdver@kam.mff.cuni.cz

[‡]Department of Mathematics, University of Ljubljana, Jadranska 19, 1111 Ljubljana, Slovenia.

sphere by inserting h crosscaps. We define *Eulerian genus* $g(S)$ of a surface S by $g(\Sigma_h) = 2h$ and $g(\Pi_h) = h$. Let $g(G)$ denote the *Eulerian genus* of the graph G , i.e., the minimal Eulerian genus of a surface into which G is embeddable.

Colorings of graphs on surfaces have been studied extensively. The fundamental result in this area is the well-known Four Color Theorem, that was proved by Appel and Haken [1] in 1977, and a shorter proof was later found by Robertson, Sanders, Seymour and Thomas [12]. Regarding the graphs on surfaces of genus $g \geq 1$, Heawood [6] showed that each graph embedded in such a surface has chromatic number at most

$$H(g) = \left\lfloor \frac{7 + \sqrt{24g + 1}}{2} \right\rfloor.$$

Later, Ringel [11] and Ringel and Youngs [10] found the corresponding lower bounds, by showing that the complete graph on $H(g)$ vertices can be embedded into any surface of Eulerian genus g , with the exception of the Klein bottle, where the correct bound on the chromatic number is 6 (established by Franklin [4]).

We consider the properties (especially regarding the chromatic number) of partitions of a graph into several subgraphs. The *partition* of a graph G to k parts consists of k edge-disjoint subgraphs G_1, \dots, G_k such that $E(G) = E(G_1) \cup E(G_2) \cup \dots \cup E(G_k)$. Note that we do not require that the subgraphs G_i are spanning, i.e., possibly $n(G_i) < n(G)$ for some i . We always assume that the graphs G_i do not contain isolated vertices. We call the subgraphs G_i *parts* of the partition.

The k -*chromatic number* $\chi_k(G)$ is the maximum of $\sum_{i=1}^k \chi(G_i)$ over all partitions G_1, G_2, \dots, G_k of G into k parts. The parameter χ_k has been studied for general graphs as well as for graphs of bounded genus. The fact that for a graph G with n vertices, $\chi_2(G) \leq n + 1$ follows from the well-known theorem of Nordhaus and Gaddum [8]. Plesník [9] proved that $n + \binom{k}{2} \leq \chi_k(K_n) \leq n + 2^{\binom{k+1}{2}}$ and conjectured that $\chi_k(K_n) = n + \binom{k}{2}$. Watkinson [15] has improved the upper bound to $\chi_k(K_n) \leq \frac{k!}{2}$ and Füredi et al. [5] to $\chi_k(K_n) \leq n + 7^k$.

Regarding the graphs with bounded genus g , let us define $\chi_k(S)$ to be maximum of $\chi_k(G)$ over all graphs G that can be embedded in S . Stiebitz and Škrekovski [14] has determined the exact values of χ_2 for all surfaces. Füredi et al. [5] have shown that

$$\chi_k(S) \leq \left\lfloor \frac{7k + \sqrt{24kg + 49k^2 - 48k}}{2} \right\rfloor,$$

and found a lower bound of order

$$\frac{7k + \sqrt{24kg + k^2}}{2}.$$

In this paper, we decrease the upper bound; this way, we obtain exact values for many surfaces and values of k .

An embedding of a graph in a surface is called *cellular* if the interior of each face is homeomorphic to an open disk. In particular, the boundary walk of each face in a cellular embedding is connected. For a face f of such an embedding, let $\ell(f)$ be the length of its boundary walk. If G is a simple connected graph with at least three vertices, then $\ell(f) \geq 3$ for each face f . A *block* of a graph G is a maximum 2-connected induced subgraph of G . Let us recall some fundamental facts about graph embeddings and surfaces that can be found e.g. in [7].

Theorem 1 *Let G be a connected graph of Eulerian genus g . Then, any embedding of G in a surface with Eulerian genus g is cellular.*

Theorem 2 (Battle et al. [2], Stahl and Beineke [13]) *If G_1, G_2, \dots, G_n are the blocks of a graph G , then*

$$g(G) = \sum_{i=1}^n g(G_i).$$

Theorem 3 (Franklin [4], Ringel [11], Ringel and Youngs [10])

Eulerian genus of the complete graph K_n is $g = \lceil \frac{1}{6}(n-3)(n-4) \rceil$. K_n can be embedded into any surface with Eulerian genus g , with the exception of K_7 , that cannot be embedded in Π_2 , i.e. the Klein bottle.

Theorem 4 (Euler's Formula) *If f is the number of faces of a cellular embedding of a graph G into a surface of Eulerian genus g , then $e(G) = n(G) + f + g - 2$.*

In the following section, we derive a version of the Euler's Formula that provides more information about a graph split into several parts (Theorem 7).

A graph G is *critical* if for every edge e of G , $\chi(G - e) < \chi(G)$. If G is a critical graph and $\chi(G) = k$, we say that G is *k-critical*. Obviously, if G is *k-critical*, then $\delta(G) \geq k - 1$. For non-complete graphs, the following stronger result known as Dirac's inequality was shown in [3]:

Theorem 5 (Dirac) *If G is a k -critical graph with $k \geq 4$ and G is not a clique, then $2e(G) \geq (k - 1)n(G) + k - 3$.*

2 Generalized Euler's Formula

Let F be the set of the faces of a cellular embedding of a simple connected graph G with at least 3 vertices. Then, $\Delta = \sum_{f \in F} (\ell(f) - 3) \geq 0$ is the number of edges that must be added to G to make it a triangulation (possibly introducing parallel edges and loops during the construction). One of the well-know consequences of Euler's Formula is the following lemma:

Proposition 6 *If G is a simple connected graph with $n \geq 3$ vertices and e edges embedded cellularly to a surface of Eulerian genus g , then $e + \Delta = 3n + 3g - 6$. In particular, $e \leq 3n + 3g(G) - 6$.*

We include the proof for the sake of completeness.

Proof. Let F be the set of faces of G . Since each edge of G appears exactly twice in the facial walks, we have $2e = \sum_{f \in F} \ell(f)$, and consequently $2e - \Delta = 3|F|$. Using Theorem 4, we infer $3e = 3n + 3|F| + 3g - 6 = 3n + 2e - \Delta + 3g - 6$, from which the desired formula immediately follows. Also, by Theorem 1, the embedding of G into a surface of Eulerian genus $g(G)$ is cellular, and since $\Delta \geq 0$, we have $e \leq 3n + 3g(G) - 6$. \square

To prove our upper bound, we need to generalize this inequality for union of several graphs:

Theorem 7 (Generalized Euler's Formula) *Let G be a simple graph and let G_1, \dots, G_k be a partition of G to k parts. Let $n_i = n(G_i) \geq 3$ for each $1 \leq i \leq k$. If every component of each G_i has at least three vertices, then*

$$e \leq 3g(G) + 3 \sum_{i=1}^k (n_i - 2).$$

Proof. Suppose that the claim is false, and let G together with its partition to graphs G_1, \dots, G_k be a counterexample that is "smallest" in the following sense:

1. $\sum_{i=1}^k (n_i - 2)$ is the smallest possible, and
2. among all graphs that satisfy the first condition, n is the largest possible.

By Proposition 6, we know that $k > 1$. Let us now describe some of the properties of G and its partition:

- (i) *Each G_i is connected.* Otherwise, we may assume without loss of generality that G_1 is not connected, i.e., $G_1 = G_1^a \cup G_1^b$, where G_1^a and G_1^b are vertex-disjoint. By the minimality, the partition $G = G_1^a \cup G_1^b \cup G_2 \cup \dots \cup G_k$ satisfies $e \leq 3g(G) + 3n(G_a) - 6 + 3n(G_b) - 6 + 3 \sum_{i=2}^k (n_i - 2) < 3g(G) + 3 \sum_{i=1}^k (n_i - 2)$, which is a contradiction with the fact that G is a counterexample.
- (ii) *G is connected.* Otherwise, G is a vertex-disjoint union of two smaller graphs G_a and G_b , and we may assume that $G_a = G_1 \cup \dots \cup G_t$ and $G_b = G_{t+1} \cup \dots \cup G_k$ (the graphs G_i are connected, thus they must be subgraphs of one of these two graphs). By Theorem 2, $g(G) = g(G_a) + g(G_b)$, and since G is a minimal counterexample, we have $e(G_a) \leq 3g(G_a) + 3 \sum_{i=1}^t (n_i - 2)$ and $e(G_b) \leq 3g(G_b) + 3 \sum_{i=t+1}^k (n_i - 2)$. Summing these two inequalities brings a contradiction with the fact that G is a counterexample.
- (iii) *Each n_i is at least 4.* Otherwise, we may assume that $n_1 = 3$ and let G' be the union of graphs G_2, \dots, G_k . Since $g(G') \leq g(G)$ and G is a minimal counterexample, it follows that $e(G') = e - e(G_1) \leq 3g(G) + 3 \sum_{i=2}^k (n_i - 2)$. However, $e(G_1) \leq 3 = 3(n_1 - 2)$, and hence $e \leq 3g(G) + 3 \sum_{i=1}^k (n_i - 2)$, which is a contradiction.
- (iv) *Minimum degree of each G_i is at least three.* Otherwise, we may assume that v is a vertex of G_1 with degree $d \leq 2$. Let $G'_1 = G_1 - v$ and let G' be the union of graphs $G'_1, G_2, G_3, \dots, G_k$. Suppose that G'_1 satisfies the assumptions of the theorem. Since $g(G') \leq g(G)$ and G is a minimal counterexample, we get $e(G') = e(G) - d \leq 3g(G) + 3 \sum_{i=1}^k (n_i - 2) - 3$, which is again a contradiction.

We need to verify that G_1 satisfies the assumptions of the theorem. This is trivial if v is not a cut-vertex of G_1 , since $n_1 \geq 4$ by the previous item. Therefore, if $d = 1$ then the assumptions are satisfied, and we may assume that G_1 does not contain a vertex of degree 1. Let us consider the case that $d = 2$ and v is a cut-vertex. Since $\delta(G_1) \geq 2$, both components of G'_1 have at least three vertices, hence in this case G'_1 satisfies the assumptions of the theorem as well.

- (v) *G is 2-connected.* Otherwise, suppose that $G = G_a \cup G_b$, where G_a and G_b share just a single vertex v . By Theorem 2, $g(G) = g(G_a) + g(G_b)$. Suppose that the graphs G_1, \dots, G_t are subgraphs of G_a , the graphs G_{t+1}, \dots, G_r are subgraphs of G_b , and for $r < i \leq k$, $G_i = G_i^a \cup G_i^b$, where G_i^a is a subgraph of G_a and G_i^b is a subgraph of G_b . Since

the minimum degree of G_i is at least three, both G_i^a and G_i^b have at least three vertices. Again, by summing the inequalities $e(G_a) \leq 3g(G_a) + 3 \sum_{i=1}^t (n_i - 2) + 3 \sum_{i=r+1}^k (n(G_i^a) - 2)$ and $e(G_b) \leq 3g(G_b) + 3 \sum_{i=t+1}^r (n_i - 2) + 3 \sum_{i=r+1}^k (n(G_i^b) - 2)$, we obtain a contradiction with the minimality of G .

- (vi) *Each two graphs G_i and G_j share at most one vertex.* Otherwise, if G_{k-1} and G_k share $t \geq 2$ vertices, then let $G'_{k-1} = G_{k-1} \cup G_k$, and apply the theorem on G split into graphs $G_1, \dots, G_{k-2}, G'_{k-1}$. We obtain $e \leq 3g(G) + 3 \sum_{i=1}^k (n_i - 2) + 6 - 3t$, which is a contradiction since $6 - 3t \leq 0$.

Let us now fix an embedding of G on a surface of Eulerian genus $g(G)$. Recall that this embedding is cellular by Theorem 1. Given a vertex v of degree d in G , let e_0, \dots, e_{d-1} be the edges of G in a cyclic ordering around v . A *segment* is a maximum interval $[a, b]$ such that all the edges e_a, e_{a+1}, \dots, e_b (with the indices taken modulo d) belong to a single graph G_i . The edges e_a and e_b are called *boundary edges* of the segment. The *length* of the segment is the number of its edges. The embedding of G has the following properties:

- *If a vertex v belongs to at least two parts, then there are at least two segments of edges at v for each of these parts.* Otherwise, suppose that all the edges of G_1 at v form just a single segment. In this case, we may split v into two vertices v_1 and v_2 such that all edges of G_1 at v are incident to v_1 and all the remaining edges at v are incident to v_2 . The created graph G' is a counterexample embedded in the same surface with $e(G') = e(G)$ and $n(G') > n(G)$, which is a contradiction with the choice of G .
- *The following configuration (\star) of edges cannot appear: $e_1 = vw$ belongs to G_i , all the remaining edges of G_i at v belong to one segment $[a, b]$, and the vertex w appears at a face f incident to e_a or e_b .* If this were the case, we might redraw G in such a way that e_1 is adjacent to e_a or e_b in the list of edges at v , by drawing it through the face f . We could then again split the vertex v , and obtain a contradiction.

We now plug the equality for Δ from Proposition 6 in the formula that we want to prove, thus obtaining the following equivalent inequality:

$$\Delta - 3n + 3 \sum_{i=1}^k n_i \geq 6k - 6.$$

Therefore, we need to show that either G has long faces, or the vertex sets of the graphs G_i have a big overlap. In fact, we prove that if the embedding of the graph G and its partition satisfies all the conditions described above, then the following stronger claim holds:

$$\Delta - 3n + 3 \sum_{i=1}^k n_i \geq 6k.$$

We proceed by the discharging method. We assign an initial charge to each vertex and each face in the following way: a vertex v that belongs to x of the graphs G_i has initial charge $3(x - 1)$. A face of length ℓ has initial charge $\ell - 3$. The sum of these charges is equal to $\Delta - 3n + 3 \sum_{i=1}^k n_i$.

Next, we move some of this charge to the graphs G_i in such a way that the final charge of each vertex and each face is nonnegative, and the final charge of each G_i is at least 6. Since no charge is lost in the process, the required inequality follows.

We use the following rules to redistribute the charge:

- (R1) Each vertex v that belongs to $x \geq 2$ graphs G_i sends charge $3/2$ to each of these graphs.
- (R2) Let f be a ≥ 4 -face and let $v_1v_2v_3v_4v_5$ be a subwalk of the facial walk of f such that edges v_2v_3 and v_3v_4 belong to the same graph G_i , and neither v_1v_2 nor v_4v_5 belongs to G_i . Then, f sends $1/2$ to G_i through each of v_2 and v_4 (one unit of charge in total).
- (R3) Let $f = wv_1v_2wv_4v_5$ be a 6-face such that the edges v_1v_2 , v_2w and v_1w belong to a graph G_i and the edges wv_4 , v_4v_5 and v_5w belong to a different graph G_j . Then, f sends $3/2$ to each of G_i and G_j through the vertex w .
- (R4) Let f be a face of length at least $t - 1$ (where $t > 5$) for that Rule R3 does not apply, and let $v_1v_2 \dots v_t$ be a subwalk of the facial walk of f such that the edges v_2v_3 , v_3v_4 , \dots , and $v_{t-2}v_{t-1}$ belong to the same graph G_i , and neither v_1v_2 nor $v_{t-1}v_t$ belongs to G_i . Then, f sends 1 to G_i through each of v_2 and v_{t-1} (two units of charge in total).

Let us first show that after the rules are applied, the final charge of each vertex and each face is nonnegative. If v is a vertex that belongs to x graphs G_i , then its final charge is zero if $x = 1$ and it is $3(x - 1) - 3x/2 = 3x/2 - 3 \geq 0$ if $x \geq 2$, by Rule R1. Now, consider the charge of the faces. Let f be an arbitrary face of G :

- (a) If Rule R3 is applied to f , then its final charge is zero.
- (b) If f is a 3-face, then either all of its edges belong to the same graph, or each of them belongs to a different graph, as otherwise two of the graphs G_i would intersect in at least two vertices. Therefore, no rule applies to f , and the final charge of f is zero.
- (c) Finally, suppose that Rule R2 applies a times and Rule R4 applies b times on an ℓ -face f . The final charge of f is $\ell - 3 - a - 2b$; therefore, it suffices to consider the case that $a + 2b + 2 \geq \ell \geq 4$. On the other hand, $\ell \geq 2a + 3b$, hence the final charge is at least $a + b - 3$, and we may assume that $a + b \leq 2$. It follows that $\ell \leq 6$ and exactly two of the graphs G_i contain edges of the face f . Since these two graphs may share only one vertex and the graph is simple, f must be a 6-face consisting of two triangles, $a = 0$ and $b = 2$. But then we obtain case (a), covered by Rule R3.

Now, let us consider the charge of the parts. We need to prove that the final charge of each of the parts is at least six. Let G_i be one of the parts, and let Y be the set of vertices that G_i shares with the rest of the graph G . Since G is 2-connected, $|Y| \geq 2$. By Rule R1, the subgraph G_i receives $3|Y|/2$ units of charge, which is at least six if $|Y| \geq 4$. Therefore, it suffices to consider the cases $|Y| = 2$ and $|Y| = 3$.

We call a boundary edge e of a segment of G_i at a vertex $v \in Y$ *rich* if e does not connect v with another vertex of Y . Let $e = vw$ be a rich edge and let f_e be a face that contains e and an edge incident to v that does not belong to G_i . Since $w \notin Y$, all the edges incident to w must belong to G_i , hence one of Rules R2, R3 or R4 applies and f_e sends at least $1/2$ units of charge through v to G_i .

Suppose first that $|Y| = 3$. Let v be an arbitrary vertex in Y . The edges of G_i at v form at least two segments. By the property (iv), the degree of v in G_i is at least 3, hence there are at least three boundary edges incident with v . Since $|Y \setminus \{v\}| = 2$, at least one of these edges is rich, hence G_i receives at least $1/2$ units of charge through v . Therefore, G_i receives $9/2$ units of charge by Rule R1, and at least $1/2$ units of charge by Rules R2–R4 through each vertex of Y , which sums to at least six units of charge.

Suppose now that $|Y| = 2$. The graph G_i receives three units of charge by Rule R1. We prove that at least $3/2$ units of charge are sent to G_i through each vertex of Y by Rules R2–R4, thus showing that G_i receives at least six units of charge. Suppose for contradiction that less than $3/2$ units of charge are sent to G_i through a vertex $v \in Y$. Then, there are at most two rich edges incident with v . On the other hand, G_i has at least two segments at v ,

the degree of v is at least three by the property (iv), and $Y \setminus \{v\}$ consists of only one vertex w , thus at least two rich edges are incident with v . Hence, we conclude that there are exactly two rich edges at v . This is only possible in the following cases:

- *The degree of v in G_i is three, and each of the edges of G_i incident with v forms a segment of length one.* However, note that in this case, each of the four (not necessarily distinct) faces incident with the rich edges sends $1/2$ units of charge through v , for total of two units of charge.
- *There are exactly two segments of G_i at v and one of them is of length one.* Let $e_0 = vu_0$ be the edge of the segment of length one, and $e_1 = vu_1$ and $e_2 = vu_2$ the boundary edges of the other segment. Note that $u_1 \neq u_2$, as the degree of v is at least three. If $w \neq u_0$ (say $w = e_1$), then each of the faces incident with e_0 send $1/2$ units of charge through v and the face f_{e_2} sends $1/2$ units of charge, for total of $3/2$ units.

Let us now consider the case that $w = u_0$. The graph G_i receives $1/2$ units of charge through v for each of e_1 and e_2 . If Rules R3 or R4 applied at v at least once, G_i would receive additional $1/2$ units of charge, contradicting the choice of v . Let us assume that this is not the case. Let w_1 and w_2 be the vertices following u_1 and u_2 in the facial walks of f_{e_1} and f_{e_2} , respectively. For $i = 1, 2$, the vertices w_i and v are both neighbours of u_i , hence $w_1 \neq v \neq w_2$. The edges following w_1 and w_2 in the facial walks do not belong to G_i , since otherwise one of Rules R3 or R4 applies. This means that $w_1, w_2 \in Y$, and hence $w_1 = w_2 = w$. This is the forbidden configuration (\star), hence we obtain a contradiction with the assumption that less than $3/2$ is sent through the vertex v .

It follows that the final charge of each of the graphs G_i is at least six, thus we conclude that $\Delta + 3(\sum_{i=1}^k n_i - n) \geq 6k$, which finishes the proof. \square

Theorem 7 is tight – for example, the equality is obtained for disjoint union of k triangulations, or graphs obtained from this graph by identifying the vertices in such a way that all edges of each graph form one segment at each vertex. Also, it is not possible to relax the condition on the number of vertices in G_i , as the claim is false if each G_i is just an edge.

3 Upper Bound

We are now ready to prove the upper bound on the k -chromatic number $\chi_k(G)$ of a graph G of Eulerian genus g . Our method is similar to the one

used by Füredi et al. [5], except that we use a better estimate on the number of edges of G obtained from Theorem 7.

Theorem 8 *Let G be a simple graph G of Eulerian genus g . If $k \leq g$, then*

$$\chi_k(G) \leq \left\lfloor \frac{7k + \sqrt{24kg + k^2}}{2} \right\rfloor.$$

Proof. Let us embed G in a surface of Eulerian genus g . Let G_1, \dots, G_k be a partition of G into k parts. For each i , let $G'_i \subseteq G_i$ be a critical subgraph of G_i such that $\chi(G'_i) = \chi(G_i) = c_i$. We may assume that $c_1 \geq c_2 \geq \dots \geq c_k$. Let t be the largest number such that $c_t \geq 7$. Thus, $t = 0$ if $c_1 \leq 6$. We bound the sum of chromatic numbers of the graphs G_1, \dots, G_t . Let $n_i = n(G'_i)$.

Let $G' = G'_1 \cup \dots \cup G'_t$ and $e' = e(G')$. Using Theorem 7, we get

$$2e' \leq 6g + 6 \sum_{i=1}^t (n_i - 2).$$

On the other hand, minimum degree of each G'_i is at least $c_i - 1$, hence $(c_i - 1)n_i \leq 2e(G'_i)$. This implies that

$$\sum_{i=1}^t (c_i - 1)n_i \leq 6g + 6 \sum_{i=1}^t (n_i - 2).$$

Using the fact that $c_i \geq 7$ and $n_i \geq c_i$, we obtain

$$\sum_{i=1}^t [(c_i - 7/2)^2 - 49/4] = \sum_{i=1}^t (c_i - 7)c_i \leq \sum_{i=1}^t (c_i - 7)n_i \leq 6g - 12t.$$

By the inequality between the arithmetic and quadratic mean,

$$\frac{1}{t} \left[\sum_{i=1}^t (c_i - 7/2) \right]^2 \leq 6g + t/4,$$

from which we infer

$$\sum_{i=1}^t c_i \leq \frac{7t + \sqrt{24tg + t^2}}{2}.$$

Taking into account the graphs G_{t+1}, \dots, G_k , we get

$$\sum_{i=1}^k c_i \leq \frac{7t + \sqrt{24tg + t^2}}{2} + 6(k - t).$$

If $t \leq g$, this expression is increasing in t , thus we obtain

$$\sum_{i=1}^k c_i \leq \frac{7k + \sqrt{24kg + k^2}}{2}.$$

Since the expression on the left-hand side is integer, we may round the expression on the right-hand side down, thus finishing the proof of this theorem. \square

4 Lower Bound

The proof of the upper bound hints at how the lower bound examples should look like. For each of the graphs in the partition, we should have $c_i = n_i$, hence all the graphs G_i should be complete. Also, since we used the inequality between arithmetic and quadratic means, their sizes should be the same. This is only possible for special values of g and k . For example, consider the case $g = \frac{1}{6}k(t - 3)(t - 4)$ for some $t \geq 4$, $t \equiv 0, 1 \pmod{3}$. Then, K_t can be embedded in a surface of genus g/k (K_7 cannot be embedded in the Klein bottle, but it can be embedded in the torus), according to Theorem 3. By Theorem 2, the disjoint union of k complete graphs on t vertices can be embedded in a surface S of genus g , hence

$$\chi_k(S) \geq kt = \frac{7k + \sqrt{24kg + k^2}}{2}.$$

For general g and k , we cannot hope for a nice formula like the one in Theorem 8, thus we would be satisfied with some description of the best possible example. A natural guess is that this example is a disjoint union of cliques. We were not able to prove that this is the case – the best result that we obtained in this direction is the following proposition:

Proposition 9 *Let G_1, \dots, G_k be a partition of a graph G of Eulerian genus g to k parts, and let $c_i = \chi(G_i) \geq 7$ for each i . Let G'_i be a c_i -critical subgraph of G_i . Suppose that $c_i \equiv 0, 1 \pmod{3}$ whenever G'_i is a clique. Then, the disjoint union of the cliques K_{c_1}, \dots, K_{c_k} has Eulerian genus at most g .*

Proof. Let $e' = e(G'_1 \cup \dots \cup G'_k)$, $n_i = n(G'_i)$, and let $\delta_i = 0$ if G'_i is a clique and $\delta_i = c_i - 3$ otherwise. By Theorem 7,

$$2e' \leq 6g + 6 \sum_{i=1}^k (n_i - 2).$$

On the other hand, using Theorem 5, we get

$$2e' \geq \sum_{i=1}^k (c_i - 1)n_i + \delta_i.$$

Therefore, we obtain

$$\begin{aligned} g &\geq \frac{1}{6} \sum_{i=1}^k (c_i - 7)n_i + 12 + \delta_i \\ &\geq \sum_{i=1}^k \frac{1}{6} ((c_i - 7)c_i + 12 + \delta_i) \\ &\geq \sum_{i=1}^k \left\lceil \frac{1}{6} (c_i - 3)(c_i - 4) \right\rceil = \sum_{i=1}^k g(K_{c_i}), \end{aligned}$$

where the last inequality holds because $\delta_i \geq 4$ whenever $c_i \equiv 2 \pmod{3}$, by the assumptions of the lemma. The statement of the lemma follows from Theorem 2. \square

5 Conclusions

Let us call the complete graph K_n *bad* if it does not triangulate the minimal surface in which it can be embedded, i.e., $n \equiv 2 \pmod{3}$. Proposition 9 shows that the best values of χ_k are achieved for disjoint unions of cliques, unless bad cliques appear in the partition. It is natural to ask whether the restriction on the appearance of the bad cliques is necessary, or whether it is always possible to “disentangle” cliques:

Problem 1 *Let G_1, \dots, G_k be a partition of a graph G to k parts such that each subgraph G_i is a clique. Is it true that the vertex-disjoint union of the cliques G_i can be embedded in a surface of Eulerian genus $g(G)$?*

For $k = 2$, this follows from Theorem 2. The proof of Theorem 7 shows that unless the graphs in the partition can be trivially disentangled, we may decrease the bound by 6, which implies that the answer to Problem 1 is positive for $k = 3$.

One way to answer the question in Problem 1 positively for $k \geq 4$ would be to improve Theorem 7, by decreasing the right hand side of the inequality by 2 for each bad clique in the partition. Another way is suggested by the following conjecture of Stiebitz and Škrekovski [14]:

Conjecture 1 *Let G be an edge-disjoint union of a clique K and an arbitrary graph H . Let H' be the graph obtained from H by contracting the set $V(K)$ to a single vertex. Then, $g(H') + g(K) \leq g(G)$.*

Because two complete graphs in a partition of a graph to k parts cannot share more than one vertex, it is easy to show by induction that Conjecture 1 implies positive answer to Problem 1.

In our considerations, we do not distinguish between orientable and non-orientable surfaces – we only focus on their Eulerian genus. While asymptotically there does not seem to be much difference, for some values k and g the results may differ.

We have provided (almost) matching upper and lower bounds for k -chromatic number of graphs with bounded genus g , assuming that the genus is large enough relatively to k . The reason why our techniques cannot be directly applied in the case k is larger than g is that we would need to consider critical graphs with chromatic number ≤ 6 . Graphs with chromatic number ≤ 4 are easy to handle – we may assume that they appear only as K_4 disjoint with the rest of the graph, since they are planar and hence do not affect genus of the graph. However, graphs with chromatic number 5 and 6 are difficult to deal with. For chromatic number 6, the list of critical graphs is known only for surfaces with $g \leq 2$, and for the chromatic number 5, there even are infinitely many of them on each surface with $g \geq 1$. Nevertheless, it might be interesting to determine the exact values of χ_k for some special cases, e.g., for graphs embedded in the torus or in the projective plane.

References

- [1] K. Appel and W. Haken, *Every planar map is four colorable*, *Contemp. Math* **98** (1989).
- [2] J. Battle, F. Harary, Y. Kodoma and J. W. T. Youngs, *Additivity of the genus of a graph*, *Bull. Amer. Math. Soc.* **68** (1962), 565–568.

- [3] G. A. Dirac, *A theorem of R. L. Brooks and a conjecture of H. Hadwiger*, Proc. London Math. Soc. (3) **7** (1957), 161–195.
- [4] P. Franklin, *A Six Colour Problem*, J. Math. Phys. **13** (1934), 363–369.
- [5] Z. Füredi, A. V. Kostochka, M. Stiebitz, R. Škrekovski and D. B. West, *Nordhaus–Gaddum-type theorems for decompositions into many parts*, J. Graph Theory **50** (2005), 273–292.
- [6] P. J. Heawood, *Map colour theorem*, Quart. J. Pure Appl. Math. **24** (1890), 332–338.
- [7] B. Mohar and C. Thomassen, *Graphs on Surfaces*, Johns Hopkins University Press, 2001.
- [8] E. A. Nordhaus and J. W. Gaddum, *On complementary graphs*, Amer. Math. Monthly **63** (1956), 175–177.
- [9] J. Plesník, *Bounds on the chromatic numbers of multiple factors of a complete graph*, J. Graph Theory **2** (1978), 9–17.
- [10] G. Ringel and J. W. T. Youngs, *Solution of the Heawood map-coloring problem*, Proc. Nat. Acad. Sci. U.S.A. **60** (1968), 438–445.
- [11] G. Ringel, *Map Color Theorem*, Springer-Verlag, New York, 1974.
- [12] N. Robertson, D. P. Sanders, P. Seymour and R. Thomas, *The four colour theorem*, J. Combin. Theory Ser. B. **70** (1997), 2–44.
- [13] S. Stahl and L. W. Beineke, *Blocks and the non-orientable genus of graphs*, J. Graph Theory **1** (1977), 75–78.
- [14] M. Stiebitz and R. Škrekovski, *A map colour theorem for the union of graphs*, J. Comb. Theory Ser. B **96** (2006), 20–37.
- [15] T. Watkinson, *A theorem of the Nordhaus–Gaddum class*, Ars Combinatoria **20-B** (1985), 35–42.