

FINITE DUALITIES, IN PARTICULAR IN FULL HOMOMORPHISMS

RICHARD N. BALL, JAROSLAV NEŠETRIL AND ALEŠ PULTR

ABSTRACT. This paper is a survey of several results concerning finite dualities, a special case of the famous Constraint Satisfaction Problem (CSP). In CSP, the point is to characterize a class \mathcal{C} of objects X determined by constraints represented by the requirement of the *existence* of structure preserving mappings *from* X into special ones. In a finite duality, such a class \mathcal{C} is characterized by the *non-existence* of special maps *into* X from a finite system of objects.

In the first third of the article we recall some well-known facts concerning constraints represented by classical homomorphisms of relational systems. In the second part we present several results, not yet published but mostly already submitted, concerning the variant of full homomorphisms. The third part contains a few results on hypergraphs and complexes in this context. These form part of an investigation recently undertaken, and appear here first.

In the Constraint Satisfaction Problem, one is concerned with objects X endowed with a given type of structure subjected to *constraints*, usually represented by a system of special objects \mathcal{B} , in our case always finite, and the requirement that there exist a mapping $X \rightarrow B \in \mathcal{B}$ suitably linked with the structures. (For a more precise formulation see Section 1 below.)

One endeavours to find a characterization, as transparent as possible, of the resulting class. This can sometimes be done by requiring the non-existence of special maps $A_i \rightarrow X$ from a finite list of objects A_1, \dots, A_n , or by requiring the non-existence of subobjects isomorphic to any of the A_i s. Then we speak of a *finite duality*.

Most of this paper is a survey. The first third is a report of some facts concerning the case of (finitary) relational systems, with the maps in question being standard homomorphisms. In the middle part we

2000 *Mathematics Subject Classification*. Primary 05E99, Secondary 05C15, 03C13, 05C15, 05C65, 08C05, 18B10.

Key words and phrases. finite model, finite structure, relational object, homomorphism, duality, hypergraph, complex.

The last two authors would like to express their thanks for support by the project 1M0021620808 of the Ministry of Education of the Czech Republic.

present several results on the case where the governing maps are full homomorphisms instead. These results have not been published yet, but are, in the majority, a part of an article that has been submitted for publication. In the last third we discuss several (mostly negative) aspects of the case where the structure is represented as a system of subsets. This is a part of a more extensive investigation that is still in its beginnings, and has not been published nor submitted yet.

1. THE CONSTRAINT SATISFACTION PROBLEM

To illustrate the type of problems and facts to be presented, let us start with the simple example of a finite binary relation. Given such relations, R on a set X and R' on a set X' , a mapping $f : X \rightarrow X'$ is a *homomorphism* $G = (X, R) \rightarrow G' = (X', R')$ if

$$(\text{hom}) \quad (x, y) \in R \Rightarrow (f(x), f(y)) \in R'.$$

Homomorphisms capture many combinatorial properties of relations; for a detailed treatment of graph homomorphisms see [9]. Of a particular interest will be the case where there is a fixed target B and we ask whether there is a homomorphism $G \rightarrow B$. That is, we are interested in the class

$$\{G \mid \text{there is an } f : G \rightarrow B\}.$$

The target B , or a system of targets \mathcal{B} , is what we speak of as a *constraint*. When we ask whether a given graph G satisfies the constraint, we are interested only in whether a homomorphism from G into B exists, and not in the homomorphism itself.

Speaking of binary relations as a *simple* example is not quite correct. Note that, already in the case of symmetric graphs and $B = K_n$ (the complete graph with n vertices), checking whether a given graph satisfies the constraint is equivalent to checking whether it is n -colorable, an extremely difficult task.

1.1. More generally, consider a category \mathfrak{C} , that is,

- a specification of *objects* of interest (relations, relational systems, hypergraphs etc.; in our case they will always be finite), and
- a specification of *morphisms*, that is maps which in one way or another respect the structure. For instance, for relational systems $(R_i)_{i \in J}$ resp. $(R'_i)_{i \in J}$ on X resp. X' , they will be the *homomorphisms* $f : (X, (R_i)) \rightarrow (X', (R'_i))$ satisfying

$$\forall i \in J, (x_1, \dots, x_{n_i}) \in R_i \implies (f(x_1), \dots, f(x_{n_i})) \in R'_i,$$

or the *full homomorphisms* $f : (X, (R_i)) \rightarrow (X', (R'_i))$ satisfying

$$\forall i \in J, (x_1, \dots, x_{n_i}) \in R_i \iff (f(x_1), \dots, f(x_{n_i})) \in R'_i.$$

The *Constraint Satisfaction Problem* (briefly, *CSP*) is that of determining for a (typically finite) system of objects \mathcal{B} the class

$$\mathbf{CSP}(\mathcal{B}) = \{X \text{ objects of } \mathfrak{C} \mid \exists B \in \mathcal{B}, \exists \text{ morphism } X \rightarrow B \text{ in } \mathfrak{C}\}.$$

2. FORBIDDING (HOMO)MORPHISMS

2.1. In a complementary way, the class $\mathbf{CSP}(\mathcal{B})$ can be represented by *forbidding* (instead of *requiring*) homomorphisms, namely as

$$\mathbf{Forb}(\mathcal{A}) = \{X \mid \text{there is no } f : A \rightarrow X \text{ with } A \in \mathcal{A}\}$$

Indeed we can take

$$(2.1) \quad \mathcal{A} = \{A \mid \text{there is no } f : A \rightarrow B \text{ with } B \in \mathcal{B}\}.$$

(If $X \in \mathbf{CSP}(\mathcal{B})$ then $X \rightarrow B \in \mathcal{B}$ and if we had $A \rightarrow X$ we would have $A \rightarrow B$; if $X \notin \mathbf{CSP}(\mathcal{B})$ then $X \notin \mathbf{Forb}(\mathcal{A})$ because of the identity $X \rightarrow X$.)

This is, of course, trivial. The less trivial question is whether we can find, for a finite \mathcal{B} , a *finite* \mathcal{A} such that

$$\mathbf{Forb}(\mathcal{A}) = \mathbf{CSP}(\mathcal{B}).$$

Then we speak of a *finite duality*. First defined in [16], finite dualities have been intensively studied from the combinatorial and logical point of view, and also in the optimization (mostly CSP) context.

Note that if one has a finite duality as above then the class $\mathbf{CSP}(\mathcal{B})$ is decidable in polynomial time. A more general (and very interesting) problem, into which we will not go here, is that of an equality $\mathbf{Forb}(\mathcal{A}) = \mathbf{CSP}(\mathcal{B})$ with at least “transparently described” \mathcal{A} , in which case we obtain a so called good characterization of $\mathbf{CSP}(\mathcal{B})$, that is, at least a deterministic decision procedure for both the positive and negative membership questions.

3. FINITE DUALITIES FOR RELATIONS WITH STANDARD HOMOMORPHISMS

The following theorem has recently been proven, as a combination of results of [2] and [18]:

3.1. Theorem. *In the case of finite binary relations, there is a finite duality $\mathbf{Forb}(\mathcal{A}) = \mathbf{CSP}(\{B\})$ if and only if the class $\mathbf{CSP}(\{B\})$ is first order definable.*

Moreover, if the finite duality exists then the \mathcal{A} can be chosen in a surprisingly special way. Namely, one has

3.2. Theorem. *If $\mathcal{B} = \{B\}$ admits a finite duality then it admits a duality $\mathbf{Forb}(\mathcal{A}) = \mathbf{CSP}(\mathcal{B})$ with \mathcal{A} a finite set of finite trees.*

For classical graphs (symmetric antireflexive relations), there are no non-trivial finite dualities. But for oriented graphs they abound ([10]), although such did not appear to be the case at the outset of the investigations. Furthermore, theorems similar to 3.1 above can be proven for relational structures of finite types; the dualities are well characterized and abundant.

3.3. Encouraged by these results, one might expect something similar for finite algebras. But the facts there are entirely different. It has been recently shown ([12]) that there are no such dualities at all. One has

Theorem. *Let Δ be a finite type. Then for every finite set \mathcal{A} of finite algebras of a type Δ and every finite algebra B of this type there exists a finite algebra A such that $A \in \mathbf{Forb}(\mathcal{A})$ and $A \notin \mathbf{CSP}(B)$.*

This not only excludes a duality, but even the existence of a finite \mathcal{A} such that

$$\mathbf{Forb}(\mathcal{A}) \subseteq \mathbf{CSP}(B).$$

Among bounded structures this is a special feature of algebras. Similar inclusions in more general relational structures may yield non-trivial classes even when there is no non-trivial duality. For instance, it can be shown that the existence of an inclusion $\mathbf{Forb}(\mathcal{A}) \subseteq \mathbf{CSP}(\mathcal{B})$ in graphs amounts to the boundedness of the chromatic numbers in $\mathbf{Forb}(\mathcal{A})$; such \mathcal{A} were characterized in [16].

For hypergraphs there is, however, a fact reminiscent of the Theorem above; see 8.3 below, and the non-existence of non-trivial inclusions $\mathbf{Forb}(\mathcal{A}) \subseteq \mathbf{CSP}(\mathcal{B})$ for complexes in 9.3.

Homomorphisms of algebras have special properties distinguishing them from general homomorphisms of relational structures. Thus for instance, for a one-one homomorphism of algebras one has

$$x = \alpha_i(x_1, \dots, x_{n_i}) \iff f(x) = \alpha'_i(f(x_1), \dots, f(x_{n_i})).$$

equivalent with the formally weaker condition with \implies . This makes them, in a sense, structurally close to full homomorphisms (recall 1.1.).

But the lack of dualities is in the specific nature of objects (algebras). As we will see below, for general relational objects the fullness condition is no obstacle to finite dualities. In fact it even helps and the dualities are much more frequent than in the standard homomorphism cases.

4. INTERMEZZO: FORBIDDEN HOMOMORPHISMS AND FORBIDDEN SUBOBJECTS

4.1. It is often the case that important classes of objects are characterized by prohibiting a *system of subobjects* rather than *homomorphisms from a system of objects*. (In fact, the idea of prohibiting subobjects emerged prior to that of prohibiting morphisms.) For example, planar graphs are characterized by the absence of two specific configurations. In the cases we have in mind it can always be done in this, perhaps more transparent, way.

Let us introduce the following notation.

$$\begin{aligned} X \rightarrow \mathcal{A} & \text{ for } \exists A \in \mathcal{A}, X \rightarrow A, \\ \mathcal{A} \rightarrow X & \text{ for } \exists A \in \mathcal{A}, A \rightarrow X, \\ X \vdash \mathcal{A} & \text{ for } \forall A \in \mathcal{A}, X \vdash A, \\ \mathcal{A} \vdash X & \text{ for } \forall A \in \mathcal{A}, A \vdash X. \end{aligned}$$

Thus, $\mathbf{Forb}(\mathcal{A}) = \mathcal{A} \vdash$, $\mathbf{CSP}(\mathcal{B}) = \rightarrow \mathcal{B}$. If we, further set

$$(4.1) \quad \mathcal{N}(\mathcal{B}) = \vdash \mathcal{B} = \{X \mid X \vdash \mathcal{B}\}$$

we see that the trivial fact from 2.1 can be expressed as

$$(4.2) \quad \mathcal{N}(\mathcal{B}) \vdash X \text{ iff } X \rightarrow \mathcal{B}$$

and the dualities we are discussing can be rewritten as

$$(4.3) \quad P \setminus (\mathcal{A} \rightarrow) = \rightarrow \mathcal{B}.$$

4.2. The categories we discuss in this paper have the following properties:

- (a) for an object A there are, up to isomorphism, only finitely many objects C such that there exists an *onto* morphism $A \rightarrow C$,
- (b) each morphism $f : X \rightarrow Y$ can be written as a composition of an onto one and an injection (that is an embedding of a subobject), symbolically

$$f = (A \twoheadrightarrow C \hookrightarrow B)$$

where we use \twoheadrightarrow to indicate morphisms onto and \hookrightarrow to indicate injections

Consequently, the duality (4.2) gives rise to the equality

$$P \setminus ((\mathcal{A} \twoheadrightarrow) \hookrightarrow) = \twoheadrightarrow \mathcal{B}.$$

and finally, setting $\mathcal{A}_1 = \mathcal{A} \twoheadrightarrow$, to

$$P \setminus (\mathcal{A}_1 \hookrightarrow) = \twoheadrightarrow \mathcal{B},$$

Hence if we write

$$\begin{aligned} \mathbf{Forb}_{\text{sub}}(\mathcal{A}_1) &= \mathcal{A}_1 \dashv \twoheadrightarrow = \\ &= \{X \mid X \text{ has no subobject isomorphic with an } A \in \mathcal{A}_1\} \end{aligned}$$

we have our original finite duality replaced by a finite “subobject duality”

$$\mathbf{Forb}_{\text{sub}}(\mathcal{A}_1) = \mathbf{CSP}(\mathcal{B}).$$

These types of situations were part of the pre-CSP motivation for studying finite dualities in [16].

5. CSP IN RELATIONAL SYSTEMS WITH FULL HOMOMORPHISMS

The categories of relational systems, and similarly the categories of hypergraphs and complexes we will be discussing later, have properties (a) and (b) from 4.2. They also obviously have the property that

(a*) every object has only finitely many subobjects.

A property that these categories do not have, but that holds true in the variant with full homomorphisms, is that

(c) each onto morphism $f : A \rightarrow B$ is a *retract*, that is, there is a $g : B \rightarrow A$ such that fg is the identity.

An object A is said to be *reduced* if it has no non-trivial (\equiv non-isomorphic) retract $r : A \rightarrow B$. Obviously, each object has a reduced retract and consequently each finite duality can be replaced with one in which all the objects in \mathcal{A} and \mathcal{B} are reduced.

An object A is *critical* with respect to a class of objects \mathcal{B} if

- it is reduced,
- $A \dashv \mathcal{B}$, and
- if $A' \rightarrow A \dashv A'$ then $A' \rightarrow \mathcal{B}$.

Set

$$\mathcal{N}_0(\mathcal{B}) = \{X \in \mathcal{N}(\mathcal{B}) \mid X \text{ critical w.r.t. } \mathcal{B}\}.$$

The following simple lemma plays a crucial role in the theorem below. What it does is reduce the $\mathcal{N}(\mathcal{B})$ from (4.1) to its essential part. The proof of the theorem will not be presented – it is in [3] – but the lemma

will be useful in a variant that does not immediately follow and that we will prove later.

5.1. Lemma. *In a category satisfying (a), (a*), (b) and (c) one has*

$$\mathcal{N}_0(\mathcal{B}) \dashv X \quad \text{iff} \quad X \rightarrow \mathcal{B}.$$

Proof. Take the $\mathcal{N}(\mathcal{B})$ from (4.1) and then consider just its reduced elements, forming a set \mathcal{A} . We still have $\mathcal{A} \dashv X$ iff $X \rightarrow \mathcal{B}$. It is easy to see that for a reduced A every morphism $A \rightarrow X$ is one-one. Thus, if $A \in \mathcal{A}$ is not critical we have a proper subobject $A' \hookrightarrow A$ such that still $A' \dashv X$. Hence we can restrict ourselves to the smallest A of \mathcal{A} (smallest in the order of “being a subobject”) and these constitute precisely the $\mathcal{N}_0(\mathcal{B})$. \square

Let Δ be any finite type. Denote by $\mathbf{Rel}_{\text{full}}(\Delta)$ the category of relational systems of this type with full homomorphisms.

5.2. Proposition. ([3]) *Let $\Delta = (n_t)_{t \in T}$ and let \mathcal{B} be a finite set of objects of $\mathbf{Rel}_{\text{full}}(\Delta)$. Let $m > \max_t n_t$. Then, with possibly finitely many exceptions, every A critical with respect to \mathcal{B} can be embedded into an object of $\mathbf{Rel}_{\text{full}}(\Delta)$ carried by X^m where*

$$X = X_B \cup \{\omega\}$$

for some $B \in \mathcal{B}$ and $\omega \notin X_B$.

As an immediate consequence one obtains

5.3. Theorem. *In $\mathbf{Rel}_{\text{full}}(\Delta)$ there exists for every finite set of objects \mathcal{B} a finite system of objects \mathcal{A} and a finite duality*

$$\mathcal{A} \dashv X \quad \text{iff} \quad X \rightarrow \mathcal{B}.$$

For graphs and one element $\mathcal{B} = \{B\}$ this was proved (a.o.) independently in [5]. Moreover, the authors have proved that one can find a duality with $|A| \leq |B| + 1$ for all the $A \in \mathcal{A}$. In this result it is essential that the graphs are not necessarily connected. For connected graphs the situation is unclear – see 7.2 below.

6. RAMSEY LISTS

In contrast with the general fact about \mathcal{B} from 5.3, it is seldom possible to complete a finite \mathcal{A} to a finite duality with \mathcal{A} on the left hand side. For instance, in the category of graphs (with full homomorphisms), there are only the following four lists \mathcal{A} with fewer than three

elements:

$$\{K_1\}, \{K_2\}, \{K_3, P_3\} \text{ and } \{K_3, P_4\}$$

(The K_n s are complete graphs, the P_n s are paths, see below). There are infinitely many such lists with three elements, however (see 7.1 below).

It is no wonder such lists are relatively rare, for they have a very strong combinatorial property.

A collection of reduced objects $\mathcal{A} = \{A_1, \dots, A_n\}$ is said to be a *Ramsey list*, or, briefly, to be *Ramsey*, if there is a finite $\mathcal{F} \subseteq \text{obj } \mathcal{C}$ such that each reduced X that is not isomorphic to an object of \mathcal{F} has a subobject isomorphic to one of the A_i s. (The reader may wish to consult [14] and [7] for a general background on Ramsey theory.)

6.1. Proposition. *Let \mathcal{C} be a category satisfying (a), (a*), (b) and (c), and let \mathcal{A} be a finite collection of reduced objects of \mathcal{C} . Then \mathcal{A} is Ramsey iff there is a finite duality*

$$\mathcal{A} \dashv X \quad \text{iff} \quad X \rightarrow \mathcal{B}.$$

Proof. If there is such a duality then it suffices to take for \mathcal{F} the set of all subobjects of the elements of \mathcal{B} .

On the other hand, if \mathcal{A} is Ramsey set $\mathcal{B} = \{F \mid \mathcal{A} \dashv F \text{ and } F \in \mathcal{F}\}$. \square

7. EXAMPLES OF CONCRETE DUALITIES IN $\mathbf{ConnGraph}_{\text{full}}$

7.1. Some particular graphs. We will use the following symbols for particular graphs. Here ij indicates that both (i, j) and (j, i) are in the relation.

- $K_n = (\{0, 1, \dots, n-1\}, \{ij \mid i \neq j\})$ is the complete graph with n vertices,
- P_n is the n -path $(\{0, 1, \dots, n\}, \{01, 12, \dots, (n-1)n\})$,
- C_n is the n -cycle $(\{0, 1, \dots, n-1\}, \{01, 12, \dots, (n-1)0\})$,
- $Y = (\{0, 1, 2, 3\}, \{01, 12, 23, 13\})$,
- $T = (\{0, 1, 2, 3, 4, 5\}, \{01, 12, 23, 34, 25\})$,
- $A = (\{0, 1, 2, 3, 4, 5\}, \{01, 12, 23, 34, 45, 14\})$,
- and $B = (\{0, 1, 2, 3, 4, 5\}, \{01, 12, 23, 34, 45, 14, 05\})$.

All the examples in this section are in the category of *connected symmetric graphs*.

7.2. Several dualities.

1. *For complete graphs we have*

$$\{K_{n+1}, P_3, Y\} \dashv X \quad \text{iff} \quad X \rightarrow K_n.$$

2. For paths:

$$\{P_4, C_3, A, C_5\} \vdash X \quad \text{iff} \quad X \rightarrow P_3,$$

and for $n \geq 4$,

$$\{P_{n+1}, T, C_3, A, B, C_5, \dots, C_{n+2}\} \vdash X \quad \text{iff} \quad X \rightarrow P_n.$$

3. For cycles:

$$\{P_4, C_3, A\} \vdash X \quad \text{iff} \quad X \rightarrow C_5,$$

and for $n \geq 6$,

$$\{P_{n-1}, T, C_3, A, B, C_5, \dots, C_{n-1}\} \vdash X \quad \text{iff} \quad X \rightarrow C_n.$$

Remarks. Note the similarities of the left duals of the paths and the cycles. Compare for instance the dualities

$$\{P_5, T, C_3, A, B, C_5, C_6\} \vdash X \quad \text{iff} \quad X \rightarrow P_4$$

and

$$\{P_6, T, C_3, A, B, C_5, C_6\} \vdash X \quad \text{iff} \quad X \rightarrow C_7.$$

7.3. A special example, and problem. By tedious checking we obtain the duality, for the A from 7.1,

$$\{P_4, C_3, C_5, E\} \vdash X \quad \text{iff} \quad X \rightarrow A$$

with

$$E = (\{0, 1, 2, 3, 4, 5, 6, 7\}, \{01, 12, 23, 34, 45, 14, 17, 26, 46, 67\}),$$

a relatively complex graph (in this context).

This contrasts with the result of [5], as $|E| = |A| + 2$, and it can be shown that the duality cannot be achieved with smaller graphs. It should not be forgotten, however, that our examples concern the category $\mathbf{ConnGraph}_{\text{full}}$, while the mentioned result speaks of general, not just connected, obstruction graphs. As far as we know, the problem of the bound on the sizes of the $A \in \mathcal{A}$ in the connected case is open.

7.4. Effectiveness of determining the left hand side.

All the examples have been established using variants of the construction from 5.2. In the case of a symmetric binary relation, the starting object does not need to be as big as X^m ; it suffices to take, roughly speaking, B redoubled with a point added, then filling in a suitable structure between the two B -layers. The search for the structure went, more or less, by brute force. Can the search be done more

effectively, if not for any graph then at least for some interesting class of graphs?

8. HYPERGRAPHS

The question naturally arises whether the general theorem 5.3 has to do with the boundedness of the type. What happens if the arity of the structure increases with the size? It turns out that, already in the simplest unbounded structure, namely in the case of hypergraphs, there are no non-trivial dualities, whether we consider the standard homomorphisms or the full ones.

8.1. A *hypergraph* is a couple $H = (V_H, \mathcal{E}_H)$ with $\mathcal{E}_H \subseteq \exp(X_H)$.

The complete hypergraph, that is any of the $S = (V_S, \mathfrak{P}(V_S))$, will be referred to as a *simplex*, for reasons that will become apparent in the next section.

A natural extension of the notion of a graph homomorphism is that of a *hypergraph homomorphism* (further, briefly, just *homomorphism*) $f : G \rightarrow H$, a mapping $f : V_G \rightarrow V_H$ such that

$$\forall X \subseteq V_G, X \in \mathcal{E}_G \implies f[X] \in \mathcal{E}_H.$$

The resulting category will be denoted by

Hypgraph.

Extending the concept of full homomorphisms from graphs to relational structures, we obtain the *full homomorphisms* between hypergraphs satisfying

$$\forall X \subseteq V_G, X \in \mathcal{E}_G \iff f[X] \in \mathcal{E}_H;$$

the resulting category will be designated

Hypgraph_{full}.

To avoid a messy discussion (caused by the fact that if $f : H \rightarrow G$ then G has a void hyperedge only when H does), we will prove the facts concerning the subcategory

Hypergraph^o

generated by the hypergraphs H such that $\emptyset \in \mathcal{E}_H$. The corresponding subcategory restricted to full homomorphisms is **Hypgraph_{full}^o**.

8.2. Proposition. *There is no non-trivial duality in **Hypergraph^o**.*

Proof. If each $B \in \mathcal{B}$ has the feature that \mathcal{E}_B contains no nonvoid edges, then each $X \in \mathbf{CSP}(\mathcal{B})$ has this feature as well. And, if \mathcal{B} were

to participate in a duality with a finite set \mathcal{A} , then no $A \in \mathcal{A}$ could have this feature. But a hypergraph X such that \mathcal{E}_X contained nonvoid edges, but such that its nonvoid edges were bigger than any of those of the \mathcal{E}_{A_S} , $A \in \mathcal{A}$, would violate the duality.

Now if there is a one-point edge in some $B \in \mathcal{B}$ then $X \rightarrow \mathcal{B}$ for any X . In this case, \mathcal{B} participates in a trivial duality with \mathcal{A} taken to be ϕ .

Finally, let there be non-void edges and let all of them have at least two points. Choose a set X such that $|X| > \max_{B \in \mathcal{B}} |V_B| \cdot \max_{A \in \mathcal{A}} |V_A|$, and set $C = (X, \mathcal{E})$, where

$$\mathcal{E} = \{M \subseteq X \mid |M| > \max_{A \in \mathcal{A}} |V_A|\}.$$

Then $C \dashv \mathcal{B}$ and $\mathcal{A} \dashv C$. \square

8.3. Recall 3.3. In $\mathbf{Hypgraph}_{\text{full}}$ we have a similar situation. (The term complex which appears in the following lemma is defined at the beginning of Section 9 below.)

Lemma. *For every system $A_1, \dots, A_k, B_1, \dots, B_l$ of hypergraphs there exists a complex C such that there is no full homomorphism $C \rightarrow B_i$ and no full homomorphism $A_j \hookrightarrow C$ unless A_j is a simplex.*

Proof. Set

$$b = \max_i |B_i|, \quad a = \max_i |A_i|.$$

Choose a set V_C with cardinality $(a+2)b$ and set

$$\mathcal{E}_C = \{E \subseteq V_C \mid |E| \leq a+1\}.$$

If f is a homomorphism into some $B \in \mathcal{B}$, choose an $x \in V_B$ such that $|f^{-1}[\{x\}]| \geq a+2$. Pick $E, E' \subseteq f^{-1}[\{x\}]$ such that $|E| = a+1$ and $|E'| = a+2$. Then $f[E] = f[E'] = \{x\}$. As $E \in \mathcal{E}_C$, $\{x\}$ is in \mathcal{E}_{B_i} . But then f is not full, since $E' \notin \mathcal{E}_C$.

Now let there be a full homomorphism $f : A_j \rightarrow C$ for some j . Let $E \subseteq V_{A_j}$ be arbitrary. Since $f[E]$ is in \mathcal{E}_C , E is in \mathcal{E}_{A_j} . \square

Corollary. *There is no non-trivial duality in $\mathbf{Hypgraph}_{\text{full}}^\circ$.*

Proof. Confront the A_i and B_j with the C from the lemma, and in addition the hypergraph D with $V_D = V_C$ and

$$\mathcal{E}_D = \{E \subseteq V_D \mid |E| = a+1 \text{ or } 0\}.$$

For the same reasons as above, $D \dashv \mathcal{B}$. There must be an $A_j \hookrightarrow D$, which makes A_j discrete, and, since $A_j \dashv B \in \mathcal{B}$, all the one-element subsets of the B 's are hyperedges. But then again the simplex among the A_i s can be mapped to any such B . \square

9. COMPLEXES

9.1. A *complex* is a hypergraph H such that

- (1) $\forall x \in V_H, \{x\} \in \mathcal{E}_H,$
- (2) $\forall E \in V_H, \forall E' \subseteq E, E' \in \mathcal{E}_H.$

(This is a well-known concept from combinatorial topology; in accordance with this we have called complete hypergraphs *simplices* – recall 8.1.)

The category of complexes with standard resp. full homomorphisms will be denoted by

Compl resp. **Compl_{full}**.

We will also be interested in the subcategory of **Compl_{full}** constituted by the complexes of dimension at most k , in other words, with the size of the hyperedges bounded by k . It will be denoted by

Compl_{full}^k.

A full homomorphism in this context is a mapping $f : G \rightarrow H$ such that

$$\forall X \subseteq V_G, |X| \leq k, \quad X \in \mathcal{E}_G \iff f[X] \in \mathcal{E}_H.$$

9.2. Proposition. *There is no non-trivial duality in **Compl**.*

Proof. Take a $B \in \mathcal{B}$ and choose an $x_0 \in V_B$. Now construct C as follows. First, let \overline{A} be the disjoint sum of all the $A \in \mathcal{A}$. Set

$$V_C = (V_B \setminus \{x_0\}) \cup V_{\overline{A}},$$

supposing the union disjoint, and define \mathcal{E}_C by

$$U \in \mathcal{E}_C \text{ if } \begin{cases} \text{either } U \in \mathcal{E}_B \text{ and } x_0 \notin U, \\ \text{or } U = (W \setminus \{x_0\}) \cup V, \quad x_0 \in W \in \mathcal{E}_B, \emptyset \neq V \in \mathcal{E}_{\overline{A}} \\ \text{or } U \in \mathcal{E}_{\overline{A}}. \end{cases}$$

Then C is a complex, $C \rightarrow B$, and $A \hookrightarrow C$ for all $A \in \mathcal{A}$. \square

9.3. Proposition. *In **Compl_{full}** there is no non-trivial finite duality. Moreover, there are no non-trivial finite sets \mathcal{A} and \mathcal{B} such that*

$$\mathbf{Forb}(\mathcal{A}) \subseteq \mathbf{CSP}(\mathcal{B}).$$

Proof. Let us specify what we understand by trivial. There is, of course, the trivial duality

$$A \dashv X \quad \text{iff} \quad X \rightarrow B$$

with B empty and A a one-point simplex; this will be excluded.

Now let \mathcal{B} contain a non-empty complex B_0 . Suppose

$$\mathbf{Forb}(\mathcal{A}) \subseteq \mathbf{CSP}(\mathcal{B}).$$

We can apply Lemma 8.3 since the C there is a complex. One of the A_j is a simplex, but then $A_j \rightarrow B_0$, which is a contradiction. \square

9.4. The proof of the following proposition needs only a very small modification of that of 5.2. But since 5.2 cannot be applied directly, we will present it in some detail. Also it is an opportunity to illustrate the principle of the proof that was omitted.

Proposition. *Let B_1, \dots, B_r be objects of $\mathbf{Compl}_{\text{full}}^k$. Let $m > k$. Then, with possibly finitely many exceptions, every A critical with respect to B_1, \dots, B_r can be embedded into an object of $\mathbf{Compl}_{\text{full}}^k$ carried by X^m where*

$$X = V_B \cup \{\omega\}$$

for some $B \in \mathcal{B}$ and $\omega \notin V_B$.

Consequently, for every finite set of objects B_1, \dots, B_r there is a finite set of objects A_1, \dots, A_n and a duality

$$\{A_1, \dots, A_n\} \dashv X \quad \text{iff} \quad X \rightarrow \{B_1, \dots, B_r\}.$$

Proof. Since A is reduced, it suffices to find a full homomorphism from A into an object as stated.

Consider a critical A . For every $a \in A$ there is a full homomorphism $i(a) : A \setminus \{a\} \rightarrow B_{i(a)}$. Assume that A is sufficiently large so as to contain distinct a_1, \dots, a_m such that the $B_{i(a_j)}$ s coincide. Designate the common value B , and choose full homomorphisms

$$f_i : A \setminus \{a_i\} \rightarrow B.$$

Set $X = V_B \cup \{\omega\}$ and define mappings

$$f_i^+ : V_A \rightarrow X$$

by setting

$$f_i^+(x) = \begin{cases} f_i(x) & \text{if } x \neq a_i, \\ \omega & \text{if } x = a_i. \end{cases}$$

Now define

$$B^+ = (X, \mathcal{E}^+) \text{ with } \mathcal{E}^+ = \{U \subseteq X \mid |U| \leq k, U \setminus \{\omega\} \in \mathcal{E}_B\}.$$

If $U \in \mathcal{E}_A$ then $f_i^+[U] \setminus \{\omega\} = f_i[U \setminus a_i] \in \mathcal{E}_B$ and hence all the f_i^+ are homomorphisms $A \rightarrow B^+$, though not necessarily full. Consider the map

$$f : A \rightarrow X^m \quad \text{defined by} \quad p_i f = f_i^+,$$

where $p_i : X^m \rightarrow X$ is the i^{th} projection. Let

$$\mathcal{E} = \{f(U) \mid U \in \mathcal{E}_A\} \cup \{U \mid |U| = 1\}$$

Thus defined, (X^m, \mathcal{E}) is an object of $\mathbf{Compl}_{\text{full}}^k$, and f and the p_i s are homomorphisms. We claim that f is full. For if $f[U] \in \mathcal{E}$ for some $U \subseteq V_A$ with $|U| \leq k$ then, since $m > k$, there is an i such that $a_i \notin U$, hence $f_i^+[U] = f_i[U]$. Therefore

$$\omega \notin f_i[U] = f_i^+[U] = p_i f[U] \in \mathcal{E}^+,$$

hence $U \in \mathcal{E}_B$, and since f_i is full, $U \in \mathcal{E}_A$. \square

9.5. It is a trivial observation that for a complex X one has

$$X \in \mathbf{Compl}_{\text{full}}^k \quad \text{iff} \quad S_{k+1} \dashv\vdash X$$

where S_{k+1} is the simplex with $k+1$ vertices. This, together with 9.4, yields the following

Corollary. *In contrast with the negative fact of 9.3, in $\mathbf{Compl}_{\text{full}}$ there exists for each finite system of objects \mathcal{B} a finite system \mathcal{A} such that*

$$\mathbf{Forb}_{\text{sub}}(\mathcal{A}) \subseteq \mathbf{CSP}(\mathcal{B}).$$

10. A FEW CONCLUDING REMARKS AND OPEN PROBLEMS

10.1. Finite dualities constitute only a small part of the CSP problem. A very important question is that of $\mathbf{Forb}(\mathcal{A})$ with \mathcal{A} not necessarily finite, but given by a criterion which is transparent. A good example is the characterization of bipartite graphs as being those into which no odd cycle embeds. Another such criterion is algorithmically generated \mathcal{A} . For such results concerning bounded tree width dualities, see [9].

10.2. In Proposition 5.2 (and similar results), the search for the elements of \mathcal{A} is restricted to subobjects of a well defined object. In some cases this suffices to present a satisfactory list, but in general the brute force search is too hard. Is there an effective search algorithm?

10.3. The existence of a non-trivial duality $\mathbf{Forb}(\mathcal{A}) = \mathbf{CSP}(\mathcal{B})$ implies the existence of a non-trivial subobject duality $\mathbf{Forb}_{\text{sub}}(\mathcal{A}) = \mathbf{CSP}(\mathcal{B})$. It would be useful to study the situations in which the latter exists and the former is absent. Note that for the ‘‘inclusion characterization’’ we have such a phenomenon in complexes: compare 9.3 with 9.5.

10.4. In the hypergraph case there are other natural choices of morphisms to be analyzed. A set of subsets can be viewed as a generalized topology, and the open continuous maps constitute one of the fullness type choices. In this particular case one has negative results similar to those in Section 8, but there are some special features of interest.

10.5. Let us recall, once again, the problem of the size of the $A \in \mathcal{A}$ bounded by the $|B|$ in the case of connected symmetric graphs with full homomorphisms (see 7.2). Does $|B| + 2$ suffice? A very easy bound is $2|B|$, obviously too big.

An analogous question for standard homomorphisms is highly non-trivial and was studied in [19].

REFERENCES

- [1] N. Alon, J. Pach, J. Solymosi, *Ramsey-type Theorems with Forbidden Subgraphs*, *Combinatorica*, **21**, **2** (2001), 155-170.
- [2] A. Atserias, *On digraph coloring problems and treewidths duality*, 20th IEEE Symposium on Logic in Computer Science (LICS) (2005), 106-115.
- [3] R.N. Ball, J. Nešetřil and A. Pultr, *Dualities in full homomorphisms*, submitted
- [4] R.N. Ball, A. Pultr and P. Vojtěchovský, *Colored graphs without colorful cycles*, to appear in *Combinatorica*.
- [5] T. Feder and P. Hell, *On Realizations of Point Determining Graphs and Obstructions to Full Homomorphisms*, to appear.
- [6] J. Foniok, J. Nešetřil, C. Tardif, *Generalized dualities and maximal finite antichains in the homomorphism order of relational structures*, to appear in *European J. Comb.*
- [7] R. L. Graham, J. Spencer, B. L. Rothschild, *Ramsey Theory*, Wiley, New York, 1980.
- [8] P. Hell, *From Graph Colouring to Constraint Satisfaction: There and Back Again*, *Topics in Discrete Mathematics*, @6. Algorithms and Combinatorics, Springer Verlag, 2006.
- [9] P. Hell, J. Nešetřil, *Graphs and Homomorphisms* Oxford University Press, Oxford, 2004.
- [10] P. Komárek, Some new good characterizations for directed graphs. *Časopis Pěst. Mat.* **109** (1984), 348–354.
- [11] G. Kun, J. Nešetřil, *Forbidden Lifts (NP and CSP for combinatorists)* submitted.
- [12] G. Kun, J. Nešetřil, *Density and Dualities for Algebras*, submitted.
- [13] S. Mac Lane, *Categories for the Working Mathematician*, Springer-Verlag, New York, 1971.
- [14] J. Nešetřil, *Ramsey Theory*, *Handbook of Combinatorics* (eds. R. L. Graham, M. Grötschel, L. Lovász), Elsevier (1995), 1331-1403.
- [15] J. Nešetřil, *Bounds and extrema for classes of graphs and finite structures. More sets, graphs and numbers*, *Bolyai soc.Math.Stud.*, 15, Springer, Berlin, 2006, 263-283.

- [16] J. Nešetřil, A. Pultr, *On classes of relations and graphs determined by subobjects and factorobjects*, Discrete Math. **22**(1978), 287-300.
- [17] J. Nešetřil, A. Pultr and C. Tardiff, *Gaps and dualities in Heyting categories*, Comment.Math.Univ.Carolinae **48,1** (2007), 9-23
- [18] J. Nešetřil, C. Tardif, *Duality theorems for finite structures (characterizing gaps and good characterizations)*, J. Comb. Th. B **80** (2000), 80-97.
- [19] J. Nešetřil, C. Tardif, *Short answers to exponentially long questions: Extremal aspects of homomorphism duality*, SIAM Journal of Discrete Mathematics **19** (2005), no. 4, 914–920.
- [20] B. Rossman, *Existential positive types and preservation under homomorphisms*, In: 20th IEEE Symposium on Logic in Computer Science (LICS),2005, pp. 467–476.

(Ball) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DENVER, DENVER, CO 80208, U.S.A.

(Nešetřil) DEPARTMENT OF APPLIED MATHEMATICS AND ITI, MFF, CHARLES UNIVERSITY, CZ 11800 PRAHA 1, MALOSTRANSKÉ NÁM. 25

(Pultr) DEPARTMENT OF APPLIED MATHEMATICS AND ITI, MFF, CHARLES UNIVERSITY, CZ 11800 PRAHA 1, MALOSTRANSKÉ NÁM. 25