A Counter–Example to Voloshin's Hypergraph Co–perfectness Conjecture

Daniel Král'
Department of Applied Mathematics and
Institute for Theoretical Computer Science (ITI)*,
Charles University,
Malostranské náměstí 25, 118 00, Prague, Czech Republic
kral@kam.mff.cuni.cz

Abstract

The upper chromatic number $\overline{\chi}(H)$ of a hypergraph H is the maximum number of colors in a coloring avoiding a polychromatic edge. The stability number $\alpha(H)$ of a hypergraph H is the cardinality of the largest set of vertices of H which does not contain an edge. A hypergraph is k–uniform if the sizes of all its edges are k. A hypergraph H is co–perfect if $\overline{\chi}(H') = \alpha(H')$ for each induced subhypergraph H' of H.

Voloshin conjectured that an r–uniform hypergraph H ($r \geq 3$) is co–perfect iff it contains neither of two special r–uniform hypergraphs (a so–called monostar and a complete circular r–uniform hypergraph on $2r-1$ vertices) as an induced subhypergraph. We disprove this conjecture for all r's.

1 Introduction

A *hypergraph* H is a pair (V, E) where V is its vertex set and $E \subseteq 2^V$ is its edge set; we do not restrict the sizes of the edges to two as in case of graphs. Throughout the paper we write $V(H)$ for a vertex set of a hypergraph H and $E(H)$ for its edge set. Recently, the topic of coloring of vertices of hypergraphs avoiding a *polychromatic* edge (i.e., the edge whose vertices have mutually different colors) has drawn an attention of different researchers, cf. [4, 6, 10, 15, 16], and related extremal (anti–Ramsey) questions were studied in [1, 3, 7, 17]. In this case, we want to color a hypergraph with a maximum possible number of colors (coloring all the vertices with the same color is clearly proper and thus minimizing the number of colors is not interesting); the maximum possible

*Institute for Theoretical Computer Science is supported by Ministry of Education of Czech Republic as project LN00A056.

number $\overline{\chi}(H)$ of colors such that the vertices of the hypergraph H can be colored avoiding a polychromatic edge is called the *upper chromatic number* of H. Besides studying this type of coloring, the researchers also study a so–called *mixed hypergraphs* where the coloring has to prevent some of the edges to be monochromatic and some of them to be polychromatic, cf. [5, 9, 11, 12, 13].

We study in this paper coloring hypergraphs which avoids a polychromatic edge as described in the previous paragraph. The *stability number* $\alpha(H)$ of a hypergraph H is the cardinality of the largest set A such that no edge of H is fully contained in A; such set A is called *stable*. If c is a coloring of the vertices of H, then a *color class* with respect to c is a set of the vertices of H colored with the same color. It is clear that $\alpha(H) \geq \overline{\chi}(H)$, since we can create a stable set by taking one vertex from each color class of a coloring using $\overline{\chi}(H)$ colors (and this is actually a stable set, since the coloring avoids a polychromatic edge). The natural question is: "For which hypergraphs H does it hold that $\alpha(H) = \overline{\chi}(H)$?" A conjecture on a possible answer to this question was stated in [18].

A hypergraph H is *r–uniform* if the sizes of all its edges are r; a hypergraph H is *r–regular* if each of its vertices is contained in precisely r edges of H. A *subhypergraph* H' of a hypergraph H is a hypergraph whose both vertex and edge sets are subsets of a vertex set and an edge set of H; a subhypergraph H' is *induced* if $E(H') = E(H) \cap 2^{V(H')}$, i.e., all the edges of H whose all the vertices are in $V(H')$ are also the edges of H'. A famous strong perfect graph conjecture asserts that $\alpha(G') = \chi(G')$ for each induced subgraph G' of G (such graphs are called *perfect*). iff G nor its complement contains an odd cycle on 5 or more vertices as an induced subgraph. Voloshin inspired by this famous conjecture made a similar conjecture in [18]:

Conjecture 1 *For each $r \geq 3$, an r–uniform hypergraph H is co–perfect if and only if it contains neither a monostar nor a C^r_{2r-1} as an induced subhypergraph.*

We postpone the missing definitions to the next paragraph. The co–perfectness of hypergraphs has been introduced in [18]; Conjecture 1 can be found as Conjecture 1 in [18]; the other conjecture stated in [18], Conjecture 2 of [18], has been recently answered in affirmative by the author in [9]. Other problems posed in [18] has been considered in [2, 8, 14] (Problem 8 of [18]), in [9] (Problem 10 and Problem 11 of [18]), in [12] (Problem 13 of [18]) and in [11, 13] (Problem 14 of [18]).

A hypergraph H is *co–perfect* if for each its induced subhypergraph H' it holds that $\alpha(H') = \overline{\chi}(H')$. A *monostar* is a hypergraph H such that the cardinality of the intersection of all the edges of H is exactly one, i.e., there exists a vertex v which is contained in all the edges and v is a unique vertex with this property; we call such a vertex the *center vertex* of a monostar. It is clear that $\alpha(H) = n-1$ for a monostar H on n vertices and $\overline{\chi}(H) < n-1$; hence monostars are certainly not co–perfect. A hypergraph H is *circular* if there exists a cycle (in the usual graph theory sense) on the vertices of H such that the edges of H form its paths; we write C^r_n for an r–uniform hypergraph whose edges are precisely all the paths consisting of r vertices of the n–vertex cycle, i.e., C^r_n is

the complete r–uniform circular hypergraph on n vertices. The hypergraph C_n^r for $n \geq 2r$ contains a monostar as an induced subhypergraph and thus it is not certainly co–perfect; but also C_{2r-1}^r is not co–perfect, since $\alpha(C_{2r-1}^r) = 2r-3$ and $\overline{\chi}(C_{2r-1}^r) < 2r-3$ (cf. [18]). These two examples of non–co–perfect hypergraphs lead to Conjecture 1 which is similar to the strong perfect graph conjecture, but besides this similarity there is no other connection between these two conjectures. Conjecture 1 has attracted attention of researchers, e.g., Tuza discussed Conjecture 1 during his invited talk at the Workshop Cycles and Colorings 2001 in Stara Lesna, Slovakia. We provide a counter–example to this conjecture for any $r \geq 3$.

Conjecture 1 is clearly equivalent to the following conjecture:

Conjecture 2 *If an r–uniform hypergraph H ($r \geq 3$) contains neither a monostar nor C_{2r-1}^r as an induced subhypergraph, then $\alpha(H) = \overline{\chi}(H)$.*

Due to Conjecture 2, it is enough to find an r–uniform hypergraph H (for each $r \geq 3$) which contains neither a monostar nor C_{2r-1}^r as an induced subhypergraph and for which $\overline{\chi}(H) < \alpha(H)$. We prove the existence of such hypergraphs in Theorem 1 in Section 2. The paper is structured as follows: We give additional definitions in Subsection 1.1. We define the counter–example r–uniform hypergraph H^r in Definition 1 of Section 2; we study properties of the hypergraph H^r in the following lemmas in Section 2 and we finish Section 2 proving Theorem 1. We conclude in Section 3.

1.1 Definitions and Used Notation

Let H be a hypergraph. We write $H \setminus V_0$ where $V_0 \subseteq V(H)$ for the induced subhypergraph of H on the vertex set $V(H) \setminus V_0$. Let c be a coloring of the vertices of H. If H contains no polychromatic edge, we say that the coloring c is *proper*. A color of a vertex v is *unique* if v is the only vertex colored with this color. An *isomorphism* between two hypergraphs H_1 and H_2 is a one–to–one mapping $\varphi : V(H_1) \rightarrow V(H_2)$ such that the images of the edges of H_1 are precisely the edges of H_2. An isomorphism is an *automorphism* if $H_1 = H_2$; an automorphism is *non–trivial* if it is not an identity. A hypergraph H is *vertex–transitive* if for any two vertices v and w of H there is an automorphism φ of H such that $\varphi(v) = w$.

The *incidence matrix* of a hypergraph H with $V(H) = \{v_1, \ldots, v_n\}$ and $E(H) = \{e_1, \ldots, e_m\}$ is $n \times m$ matrix $I(H)$ such that $I(H)_{ij} = 1$ if $v_i \in e_j$ and $I(H)_{ij} = 0$ otherwise. Note that if H is r–uniform, then each column sum is precisely r; if H is k–regular, then each row sum is precisely k. We deal with different uniform hypergraphs in the paper: We try to use the notation such that the superscript is equal to the common sizes of edges, e.g., C_n^r (defined earlier) is an r–uniform hypergraph.

2 The Counter–Example

We first define the counter–example (to Conjecture 2) r–uniform hypergraph H^r:

Definition 1 *Let $r \geq 3$ be a fixed integer. Let H^r be the r–uniform hypergraph with $2r$ vertices and $2r+2$ edges whose incidence matrix $2r \times (2r+2)$ is the following (the incidence matrices for $r=3$ and $r=4$ can be found below):*

$$I(H^r) = \left(\begin{array}{ccccccccccc|cc} 1 & 0 & 0 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 & 1 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 & 1 & 1 & 0 & 1 \end{array}\right)$$

We write $v_1, \ldots, v_{2r}$ for the vertices of H^r; the vertex v_i corresponds to the i–th row of the incidence matrix. We write $e_1, \ldots, e_{2r}$ for the edges corresponding to the first $2r$ columns of the incidence matrix; the edge e_i corresponds to the i–th column of the incidence matrix:

$$e_i = \{v_i, v_{i+1}, v_{i+3}, \ldots, v_{i+2r-3}\}$$

where the subscripts of the vertices are taken modulo $2r$. We write e_o and e_e (odd and even in correspondence to the parity of the indices of the subscripts of the vertices contained in e_o and e_e) for the edges corresponding to the last but one and the last column of the incidence matrix.

In order to illustrate the definition, we include the incidence matrices for H^3 and H^4:

$$I(H^3) = \left(\begin{array}{cccccc|cc} 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \end{array}\right)$$

$$I(H^4) = \left(\begin{array}{cccccccc|cc} 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \end{array}\right)$$

Lemma 1 *The hypergraph H^r is a vertex–transitive r–uniform $(r+1)$–regular hypergraph.*

Proof: The proof of the uniformity and the regularity of H^r follows immediately from Definition 1. In order to prove the vertex–transitivity of H^r, note that the function $\varphi : V(H^r) \rightarrow V(H^r)$ defined as follows is an automorphism of H^r:

$$\varphi(v_i) = \begin{cases} v_1 & \text{if } i = 2r, \\ v_{i+1} & \text{otherwise.} \end{cases}$$

■

We next find the stability number of H^r:

Lemma 2 *The stability number of H^r is $2r-3$.*

Proof: Let $r \geq 3$ be a fixed integer through the proof. The set of vertices of H^r $\{v_1, \ldots, v_{2r-3}\}$ is stable; thus $\alpha(H^r) \geq 2r-3$. If $\alpha(H^r) > 2r-3$, let $A \subseteq V(H^r)$ be the stable set of size $2r-2$. We can assume that $v_1 \notin A$ since H^r is vertex–transitive. Let v_i be the only vertex different from v_1 not contained in A. If i is odd, then $e_e \subseteq A$. Hence i has to be even. If $i = 2$, then $e_3 \subseteq A$; but if $i \geq 4$, then $e_2 \subseteq A$.

■

We next prove a lemma which allows us to use a proper coloring of H^r to get a proper coloring of H^{r-1}:

Lemma 3 *Let c be a coloring of the hypergraph $H^r, r \geq 4$, using k colors such that the colors of v_{2r-1} and v_{2r} are unique. Then $\overline{\chi}(H^{r-1}) \geq k-2$.*

Proof: We write e'_i for the i–th edge of H^{r-1} and e_i for the i–th edge of H^r in the proof; we consider the vertices $v_i, 1 \leq i \leq 2r-2$ of H^{r-1} and H^r to be the same ones; we write V_0 for these vertices, i.e., $V_0 = \{v_1, v_2, \ldots, v_{2r-2}\}$. Let c' be the coloring of H^{r-1} obtained by restriction of c to the first $2r-2$ vertices, i.e., $c'(v_i) = c(v_i), 1 \leq i \leq 2r-2$. We claim that there is no polychromatic edge in H^{r-1}. The edge e'_e, resp. e'_o, is not polychromatic, since $e'_e = e_e \cap V_0$, resp. $e'_o = e_o \cap V_0$ (recall that the colors of the vertices v_{2r-1} and v_{2r} are different and not used to color any of the vertices of V_0). The edges $e'_i, 2 \leq i \leq 2r-2$, are not

polychromatic, since $e'_i = e_i \cap V_0$. The remaining edge e'_1 is not polychromatic, since $e'_1 \subset e_{2r-1} \cap V_0$.

■

We next prove that the upper chromatic number of H^r is smaller than its stability number:

Lemma 4 *The upper chromatic number of H^r is $2r-4$.*

Proof: Let c be the following coloring of H^r:

$$c(v_i) = \begin{cases} i & \text{for } 1 \leq i \leq 2r-4, \\ 2r-4 & \text{for } 2r-3 \leq i \leq i. \end{cases}$$

The coloring c is a proper coloring of H^r and thus $\overline{\chi}(H^r) \geq 2r-4$. In the rest, we prove that $\overline{\chi}(H^r) \leq 2r-4$; the proof proceeds by induction on r (although it might not seem so in the beginning of it):

- $r = 3$
  Let c be a coloring of H^3 using 3 colors. The coloring c colors some consecutive vertices by different colors and hence we may assume that $c(v_1) \neq c(v_2)$ due to the vertex–transitivity of H^3. We distinguish three cases according to the possibilities which vertex is assigned the third color (the one different from $c(v_1)$ and $c(v_2)$):

  - The third color is assigned to v_3. (This case is symmetric to the case that the third color is assigned to v_6).
    The color $c(v_5)$ is $c(v_3)$ due to the edges $e_2 = \{v_2, v_3, v_5\}$ and $e_o = \{v_1, v_3, v_5\}$. The color $c(v_6)$ is $c(v_3)$ due to the edges $e_5 = \{v_5, v_6, v_2\}$ and $e_6 = \{v_6, v_1, v_3\}$. Then, $c(v_4)$ has to be $c(v_2)$ due to the edges $e_1 = \{v_1, v_2, v_4\}$ and $e_e = \{v_2, v_4, v_6\}$. But then the edge $e_4 = \{v_4, v_5, v_1\}$ is polychromatic — contradiction.
  - The third color is assigned to v_4.
    This is impossible, since $e_1 = \{v_1, v_2, v_4\}$.
  - The third color is assigned to v_5.
    Then $c(v_3) \in \{c(v_2), c(v_5)\}$ due to the edge $e_2 = \{v_2, v_3, v_5\}$ and $c(v_3) \in \{c(v_1), c(v_5)\}$ due to the edge $e_o = \{v_1, v_3, v_5\}$. Hence $c(v_3) = c(v_5)$, the third color is assigned to v_3 and this case has been dealt in the first subcase.

- $r = 4$
  Let c be a coloring of H^4 using 5 colors; we distinguish several cases according to the sizes of color classes of c:

  - $4:1:1:1:1$
    If there were two consecutive (with respect to their subscripts) vertices with unique colors, Lemma 3 would imply that $\overline{\chi}(H^3) \geq 3$.

Since there are four vertices with unique colors, we can assume w.l.o.g. that they are v_1, v_3, v_5, v_7 (no two of them can be consecutive and H^4 is vertex–transitive). But then edge e_o is polychromatic — contradiction.

- $3:2:1:1:1$
  Let C_1 be the color class of size 3 and C_2 be the color class of size 2. Since it cannot hold both $|C_1 \cap e_o| \geq 2$ and $|C_1 \cap e_e| \geq 2$, we may assume that $C_2 \subseteq e_e$. Then $|C_1 \cap e_o| \geq 2$. We assume that $v_1 \in C_1$ (recall that H^4 is vertex–transitive). We prove that $C_1 \subseteq \{v_1, v_2, v_3, v_5, v_7\}$. Consider the edge $e_2 = \{v_2, v_3, v_5, v_7\}$. Since $|e_2 \cap C_2| \leq 1$, it has to be $|e_2 \cap C_1| = 2$. Hence $C_1 \subseteq \{v_1, v_2, v_3, v_5, v_7\}$. Let v_i be one of the two vertices of C_2; note that $i \in \{2, 4, 6, 8\}$. We assume that $i \neq 8$ (otherwise take the other vertex of C_2). Consider the edge e_{i+1}: $|e_{i+1} \cap C_2| \leq 1$ and thus it has to be $|e_{i+1} \cap C_1| \geq 2$ and $C_1 = \{v_1, v_2, v_{i+1}\}$. Then the edge e_{i+2} is polychromatic ($e_{i+2} \cap C_1 \subseteq \{v_1\}$ and $e_{i+2} \cap C_2 \subseteq \{v_{i+2}\}$) — contradiction.
- $2:2:2:1:1$
  Let C_1, C_2 and C_3 be the three color classes of sizes 2. We assume w.l.o.g. that $C_2 \subset e_o$ and $C_3 \subset e_e$ (otherwise, the edge e_o or the edge e_e is polychromatic). Let v_i be a vertex of either C_2 or C_3. Hence $|e_{i+1} \cap C_2| \leq 1$ and $|e_{i+1} \cap C_3| \leq 1$; this gives that $C_1 \subseteq e_{i+1}$ (the subscript is considered modulo 8). But then C_1 has to be in the intersection of four different edges $e_{k_1}, e_{k_2}, e_{l_1}, e_{l_2}$ where k_1 and k_2 are odd and l_1 and l_2 are even; but any such intersection is empty, since $e_{k_1} \cap e_{k_2} \subseteq \{v_2, v_4, v_6, v_8\}$ and $e_{l_1} \cap e_{l_2} \subseteq \{v_1, v_3, v_5, v_7\}$ — contradiction.

- $r = 5$
  Let c be a coloring of H^5 using 7 colors; we distinguish several cases according to the sizes of color classes of c:

  - $4:1:1:1:1:1:1$
    In this case, there are certainly two consecutive (with respect to their subscripts) vertices with unique colors, w.l.o.g. v_9 and v_{10}. But then Lemma 3 would imply that $\overline{\chi}(H^4) \geq 5$ — contradiction.
  - $3:2:1:1:1:1:1$
    There cannot be two consecutive vertices with unique colors as dealt in the previous subcase. But then w.l.o.g. the colors of v_1, v_3, v_5, v_7, v_9 have to be unique (there are five vertices with unique colors and H^5 is vertex–transitive) and the edge e_o is polychromatic — contradiction.
  - $2:2:2:1:1:1:1$
    Let C_1, C_2 and C_3 be the three color classes of sizes 2. We assume w.l.o.g. that $C_2 \subset e_o$ and $C_3 \subset e_e$ (otherwise, these edges would be polychromatic). Let v_i be a vertex of either C_2 or C_3. Hence

$|e_{i+1} \cap C_2| \leq 1$ and $|e_{i+1} \cap C_3| \leq 1$; this implies that $C_1 \subseteq e_{i+1}$ (the subscript is considered modulo 10). But then C_1 has to be in the intersection of four different edges $e_{k_1}, e_{k_2}, e_{l_1}, e_{l_2}$ where k_1 and k_2 are odd and l_1 and l_2 are even; but any such intersection is empty, since $e_{k_1} \cap e_{k_2} \subseteq \{v_2, v_4, v_6, v_8, v_{10}\}$ and $e_{l_1} \cap e_{l_2} \subseteq \{v_1, v_3, v_5, v_7, v_9\}$ — contradiction.

- $r = 6$
  Let c be a coloring of H^6 using 9 colors; H^6 contains at least 6 vertices with unique colors. If there were two consecutive (with respect to their subscripts) vertices with unique colors, we may assume that these are v_{11} and v_{12} since H^6 is vertex–transitive, Lemma 3 would imply that $\overline{\chi}(H^5) \geq 7$. If there were more than 6 vertices with unique colors, then (due to pigeonhole principle) there would be two consecutive ones. Thus there are exactly 6 vertices colored with unique colors and these are w.l.o.g. $v_1, v_3, v_5, v_7, v_9, v_{11}$ (no two of them can be consecutive and H^6 is vertex–transitive). But then edge e_o is polychromatic — contradiction.

- $r \geq 7$
  Let c be a coloring of H^r using $2r-3$ colors. H^r contains at least $2r-6 > r$ vertices with unique colors. Due to pigeonhole principle, there are two consecutive (with respect to their subscripts) vertices colored with unique colors. We may assume that these vertices are v_{2r-1} and v_{2r}, since H^r is vertex–transitive, but then Lemma 3 implies that $2r - 5 \leq \overline{\chi}(H^{r-1})$ — contradiction.

■

It remains to check that H^r contains neither a monostar nor C^r_{2r-1} as an induced subhypergraph:

Lemma 5 *The hypergraph H^r does not contain a monostar as an induced subhypergraph.*

Proof: Let $r \geq 3$ be a fixed integer through the proof. We assume that H^r contains a monostar with the center vertex equal to v_1. Let $V_0 \subseteq V(H^r)$ be the vertices which induce the monostar and let $E_0 = E(H^r) \cap 2^{V_0}$. Note that the following hold due to the definition of a monostar and an induced subhypergraph:

$$V_0 = \bigcup_{e \in E_0} e$$

$$\forall e' \in E(H) : e' \subseteq V_0 \Rightarrow e' \in E_0$$

$$\{v_1\} = \bigcap_{e \in E_0} e = \bigcap_{e \subseteq V_0, e \in E(H')} e$$

We distinguish several cases in the proof:

- $e_1 \in E_0$ and $e_o \in E_0$
  It has to be that $V_0 \supseteq e_o \cup e_1 = V(H^r) \setminus \{v_{2r}\}$. But then $e_2 \in E_0$ — contradiction.

- $e_1 \in E_0$ and $e_o \notin E_0$
  The edge e_1 cannot be the only edge of E_0. Because the intersection of the edges of E_0 is $\{v_1\}$, E_0 can contain besides e_1 only the edges e_i for even $i, 4 \leq i \leq 2r$. If $e_{2r} \in E_0$, then $e_1 \cup e_{2r} = V(H^r) \setminus \{v_{2r-1}\} \supseteq V_0$ and $e_e \in E_0$ which is impossible. Let e_{i_1} be an edge of E_0 different from e_1; i_1 has to be an even integer between 4 and $2r-2$. Since $e_1 \cap e_{i_1} = \{v_1, v_{i_1}\}$, the edge set E_0 has to contain an edge e_{i_2} different from e_1 and e_{i_1}. But then $e_1 \cup e_{i_1} \cup e_{i_2} = V(H^r) \setminus \{v_{2r}\} \supseteq V_0$. Hence $e_2 \in E_0$ — contradiction.

- $e_1 \notin E_0$ and $e_o \in E_0$
  The only two edges of E_0 which contain v_1 and do not contain v_i for odd $3 \leq i \leq 2r-1$ are e_1 and e_{i+1}. Since $e_1 \notin E_0$, the intersection of the edges of E_0 consists of the single vertex v_1 and $v_i \in e_o$ for all odd i's, $3 \leq i \leq 2r-1$, it has to be $e_{i+1} \in E_0$. But then, $V_0 \supseteq V(H^r) \setminus \{v_2\}$ and e_3 has to be contained in E_0 — contradiction.

- $e_1 \notin E_0$ and $e_o \notin E_0$
  In this case it has to be $E_0 \subseteq \{e_4, e_6, \ldots, e_{2r}\}$. The only edge of e_4, e_6, ..., e_{2r} which does not contain v_i for odd $i, 3 \leq i \leq 2r-1$ is e_{i+1}. Hence, it has to be $E_0 = \{e_4, \ldots, e_{2r}\}$. But then, $V_0 \supseteq V(H^r) \setminus \{v_2\}$ and thus e_3 has to be contained in E_0 — contradiction.

■

Lemma 6 *The hypergraph H^r does not contain the complete circular hypergraph C^r_{2r-1} on $2r-1$ vertices as an induced subhypergraph.*

Proof: Let $r \geq 3$ be a fixed integer through the proof. If H^r contains C^r_{2r-1} as an induced subhypergraph, then $H^r \setminus v_1$ is isomorphic to C^r_{2r-1} (recall that H^r is vertex–transitive). But $H^r \setminus v_1$ consists of only $2r + 2 - (r+1) = r+1$ edges and C^r_{2r-1} consists of $2r-1$ edges.

■

We can conclude the section:

Theorem 1 *The r–uniform hypergraph H^r contains neither a monostar nor the complete circular hypergraph C^r_{2r-1} on $2r-1$ vertices, but $\overline{\chi}(H^r) < \alpha(H^r)$ for any $r \geq 3$.*

Proof: The proof immediately follows from Lemma 2, Lemma 4, Lemma 5 and Lemma 6.

■

3 Conclusion

The co–perfectness conjecture of Voloshin is wrong, but the following interesting problem remains open:

Problem 1 *For which (r–uniform) hypergraphs H does the equality $\alpha(H) = \overline{\chi}(H)$ hold?*

The answer suggested by Conjecture 2 for r–uniform hypergraphs is false. We do not think there is some hope to find a finite number of cases of minimal non–perfect hypergraphs different from monostars, but even though the answer to Problem 1 could be nice and provide new insights to the structure of colorings avoiding polychromatic edges.

Acknowledgement

The author would like to thank Jan Kára for reading an early version of this manuscript and pointing out an error which it contained. The author would also like to thank Jan Kratochvíl for introducing the concept of mixed hypergraphs and discussions on the topic.

References

[1] N. Alon: On a conjecture of Erdös, Simonovits, and Sós concerning anti–Ramsey theorems, *J. Graph Theory* **1**, 1983, 91–94.

[2] Z. Dvořák, D. Král': On Planar Mixed Hypergraphs, *Electronic J. Combin.* **8 (1)**, 2001, #R35.

[3] P. Erdös, M. Simonovits, V. T. Sós: Anti–Ramsey theorems, *Coll. Math. Soc. J. Bolyai* **10**, Infinite and finite sets, Keszthely, Hungary, 1973, 657–665.

[4] T. Jiang: Edge–colorings with no large polychromatic stars, to appear in *Graphs and Combinatorics.*

[5] T. Jiang, D. Mubayi, Zs. Tuza, V. Voloshin and D. B. West: Chromatic spectrum is broken, to appear in *Graphs and Combinatorics.*

[6] T. Jiang, D. B. West: Edge–Colorings of Complete Graphs that Avoid Polychromatic Trees, to appear in *Discrete Mathematics.*

[7] T. Jiang, D. B. West: On the Erdos–Simonovits–Sos Conjecture on the anti–Ramsey number of a cycle, submitted.

[8] D. Kobler, A. Kündgen: Gaps in the chromatic spectrum of face–constrained plane graphs, *Electronic J. Combin.* **3**, 2001, #N3.

[9] D. Král': On Feasible Sets of Mixed Hypergraphs, submitted.

[10] D. Král': On Maximum Face–Constrained Coloring of Plane Graphs of Girth at least 5, submitted.

[11] D. Král', J. Kratochvíl, H.-J. Voss: Mixed Hypercacti, submitted.

[12] D. Král', J. Kratochvíl, H.-J. Voss: Mixed Hypergraphs with Bounded Degree: Edge–Colouring of Mixed Multigraphs, to appear in *Theoretical Computer Science*.

[13] D. Král', J. Kratochvíl, H.-J. Voss: Mixed Hypertrees, submitted.

[14] A. Kündgen, E. Mendelsohn, V. Voloshin: Colouring planar mixed hypergraphs, *Electronic J. Combin.* **7**, 2000, #R60.

[15] A.Kündgen, R. Ramamurthi: Coloring face–hypergraphs of graphs on surfaces, to appear in *Journal of Combinatorial Theory (B)*.

[16] R. Ramamurthi, D. B. West: Maximum Face–Constrained Coloring of Plane Graphs, to appear in *Discrete Mathematics*.

[17] M. Simonovits, V. T. Sós: On restricting colorings of K_n, *Combinatorica* **4**, 1984, 101–110.

[18] V. Voloshin, On the upper chromatic number of a hypergraph, *Australasian Journal of Combinatorics* **11**, 1995, 25–45.