

# Non-intersecting perfect matchings in cubic graphs

Tomáš Kaiser<sup>1</sup>

André Raspaud<sup>2</sup>

## Abstract

A conjecture of G. Fan and A. Raspaud asserts that every bridgeless cubic graph contains three perfect matchings with empty intersection. We suggest a possible approach to problems of this type, based on the concept of a balanced join in an embedded graph. We use this method to prove a special case of a conjecture of E. Máčajová and M. Škoviča on Fano colorings of cubic graphs.

## 1 Introduction

A number of problems involving cubic graphs concerns the existence of perfect matchings whose intersection is small or empty. This is natural as the existence of two disjoint perfect matchings in a cubic graph  $G$  is equivalent to  $G$  being 3-edge-colorable, a fundamental property in the world of cubic graphs. Although there are cubic graphs (even without bridges) that do not have this property, the following has been conjectured by Fan and the second author [2] in 1994:

**Conjecture 1.** *Every bridgeless cubic graph contains perfect matchings  $M_1$ ,  $M_2$ ,  $M_3$  such that*

$$M_1 \cap M_2 \cap M_3 = \emptyset.$$

(The definitions of all the notions used in this introductory section will be found in Section 2.)

We remark that Conjecture 1 would be implied by the celebrated Berge–Fulkerson conjecture [4] (see also [9]). Another related set of problems, studied, e.g., in [6, 7], concerns so-called Fano colorings. Let  $G$  be a cubic graph. A *Fano*

---

<sup>1</sup>Department of Mathematics and Institute for Theoretical Computer Science, University of West Bohemia, Univerzitní 8, 306 14 Plzeň, Czech Republic. E-mail: [kaisert@kma.zcu.cz](mailto:kaisert@kma.zcu.cz). Supported by project 1M0545 and Research Plan MSM 4977751301 of the Czech Ministry of Education. The research was partly conducted while the first author was visiting LaBRI; the visit was supported by the COMBSTRU network.

<sup>2</sup>LaBRI, Université Bordeaux I, 351, cours de la Libération, 33405 Talence cedex, France. E-mail: [raspaud@labri.fr](mailto:raspaud@labri.fr).




Figure 1: The Fano plane  $\mathcal{F}_7$ .

*coloring* of  $G$  is any assignment of points of the Fano plane  $\mathcal{F}_7$  (see Figure 1) to edges of  $G$  such that the three edges incident with each vertex of  $G$  are mapped to three distinct collinear points of  $\mathcal{F}_7$ . For  $k \leq 7$ , a *k-line Fano coloring* is a Fano coloring in which only at most  $k$  lines of  $\mathcal{F}_7$  appear as color patterns at the vertices.

It is shown in [7] that every bridgeless cubic graph admits a 6-line Fano coloring. Furthermore, bridgeless cubic graphs are conjectured to admit 4-line Fano colorings, and it is observed that the conjecture is equivalent to Conjecture 1. A natural intermediate conjecture on 5-line Fano colorings is stated in [7] as Conjecture 5.1; we give it in a slightly less specific form:

**Conjecture 2.** *Every bridgeless cubic graph admits a 5-line Fano coloring.*

We will refer to this conjecture as the *5-line conjecture*. Before we present a (more or less well-known) equivalent formulation in the spirit of Conjecture 1, we need an important definition. A *join* in a graph  $G$  is a set  $J \subset E(G)$  such that the degree of every vertex in  $G$  has the same parity as its degree in the graph  $(V(G), J)$ . (In the literature, the terms *postman join* or *parity subgraph* have essentially the same meaning.)

It can be shown that the following is equivalent to Conjecture 2:

**Conjecture 3.** *Every bridgeless cubic graph admits two perfect matchings  $M_1$ ,  $M_2$  and a join  $J$  such that*

$$M_1 \cap M_2 \cap J = \emptyset.$$

In the present paper, we suggest a possible approach to problems like Conjectures 1 and 3, based on what we call ‘balanced joins’ in embedded graphs (see Section 3). The technique was inspired by that of [7] and may be regarded as its refinement. In Section 4, we relate balanced joins in an embedded graph to independent sets in its dual.

Our main result is a special case of the 5-line conjecture. Recall that the *oddness* of a cubic graph  $G$  is the minimum number of odd circuits in a 2-factor of  $G$ . In Section 6 we prove:

**Theorem 4.** *Conjecture 2 (and Conjecture 3) is true for bridgeless cubic graphs of oddness two.*

A major tool in the proof of Theorem 4 is a splitting lemma (Lemma 13) proved in Section 5. In the concluding section, we give several intriguing open problems.

## 2 Preliminaries

Our graphs may contain loops and parallel edges. The vertex and edge sets of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$ , respectively. Each edge is viewed as composed of two *half-edges* (that are *associated* to each other) and we let  $E(v)$  denote the set of half-edges incident with a vertex  $v$ . To keep notation simple, we do not distinguish between a set of edges  $F \subset E(G)$  and the set of half-edges composing the edges in  $F$ .

The *degree* of a vertex  $v \in V(G)$  is denoted by  $d_G(v)$ . A loop at  $v$  contributes 2 to  $d_G(v)$ . For  $X \subset V(G)$ , we let  $\partial_G(X)$  be the set of edges with exactly one endvertex in  $X$ .

As usual, e.g., in the theory of nowhere-zero flows, we define a *cycle* in a graph  $G$  to be any subgraph  $H \subset G$  such that each vertex of  $H$  has even degree in  $H$ . (Thus, a cycle need not be connected.)

**Observation 5.** *A subgraph  $H \subset G$  is a cycle in  $G$  if and only if  $E(G) - E(H)$  is a join.*

An *edge-cut* (or just *cut*) in  $G$  is a set  $C \subset E(G)$  such that  $G - C$  has more components than  $G$ , and  $C$  is inclusionwise minimal with this property. A *bridge* is a cut of size 1. A graph is *bridgeless* if it contains no bridge (note that with this definition, a bridgeless graph may be disconnected). A  *$k$ -edge-connected* graph (where  $k \geq 1$ ) is one that is connected and contains no cut of size at most  $k - 1$ . All these notions naturally extend to half-edges; for instance, we may say that a set of half-edges forms a cut in  $G$ .

A cut  $C$  in  $G$  is *odd* if  $|C|$  is odd. The *odd edge-connectivity*  $\lambda^{\text{odd}}(G)$  of  $G$  is the size of a smallest odd cut in  $G$ , or  $\infty$  if no such cut exists.

For terms not defined here, we refer the reader to any standard textbook of graph theory such as [1].

## 3 Balanced joins in embedded graphs

We begin by recalling a combinatorial representation of graphs embedded on surfaces, due to Heffter, Edmonds and Ringel (see, e.g., [8, Chapter 3]). A *rotation system* for  $H$  is a mapping assigning to every vertex  $v \in V(H)$  a cyclic ordering of the half-edges in  $E(v)$  (a *rotation at  $v$* ). A rotation system for  $H$  corresponds to an embedding of  $H$  in some orientable surface. Accordingly, we will speak of a graph  $H$  with a given rotation system as an *embedded graph*. For half-edges  $h, h'$




Figure 2: (a) The Petersen graph  $\mathcal{P}$  with a perfect matching  $M$  (bold). (b) An  $M$ -contraction  $\tilde{\mathcal{P}}$  of  $\mathcal{P}$ ; the clockwise ordering of half-edges at vertices specifies the rotation system.

incident with a vertex  $v$ , we may say that  $h$  *precedes* or *follows*  $h'$  at  $v$ , or that  $h$  and  $h'$  are *consecutive* at  $v$ , always referring to the rotation order.

Cubic graphs and embedded graphs are related in the following way. Let  $G$  be a cubic graph and  $M$  a perfect matching in  $G$ , and let  $o$  be an orientation of the complementary 2-factor  $\overline{M}$ . Contract each circuit  $C$  of  $\overline{M}$  to a vertex  $v_C$ . The resulting graph  $G_{M,o}$  is an embedded graph: the rotation at  $v_C$  is given by the order in which the corresponding half-edges meet the circuit  $C$  in  $G$ . A graph of the form  $G_{M,o}$  for some orientation  $o$  will be called an  $M$ -*contraction* of  $G$ . We identify the edges of  $G_{M,o}$  with the corresponding edges of  $G$ .

As an example, consider the Petersen graph  $\mathcal{P}$  and the perfect matching  $M$  shown in Figure 2a. The result,  $\tilde{\mathcal{P}}$ , of the above process (for a suitable orientation  $o$ ) is shown in Figure 2b. Note that we get essentially the same embedded graph for any other choice of  $M$  and  $o$ . An embedding of  $\tilde{\mathcal{P}}$  corresponding to its rotation system (in an orientable surface of genus 3) is shown in Figure 3.

Let  $H$  be an arbitrary embedded graph. For  $z \in V(H)$  and  $h, h' \in E(z)$ , we define  $[h, h']$  as the set of half-edges at  $z$  containing  $h, h'$  and all the half-edges encountered as one passes from  $h$  to  $h'$  in the direction given by the rotation at  $v$ . Note that if  $h = h'$ , then  $[h, h'] = E(z)$ .

For  $J \subset E(H)$ , a  $J$ -*segment* (at  $z$ ) is any set of the form  $[h, h']$ , where  $h$  and  $h'$  are half-edges incident with  $z$  and  $[h, h'] \cap J = \{h, h'\}$ . In addition, if  $J \cap E(z) = \emptyset$ , we consider  $E(z)$  as a  $J$ -segment. The *length* or *size* of a  $J$ -segment is its cardinality as a set.

We define a *balanced join* in  $H$  to be a set  $J \subset E(H)$  such that for each vertex  $v$ , every  $J$ -segment at  $v$  has even length. The choice of the term is partly justified by the following observation:

**Observation 6.** *Every balanced join in an embedded graph is a join.*




Figure 3: An embedding of the graph  $\tilde{\mathcal{P}}$  of Figure 2b in an orientable surface of genus 3 determined by its rotation system. The opposite edges of the bounding octagon are identified (in such a way that the arrows match).

*Proof.* Let  $J$  be a balanced join in  $H$  and  $v \in V(H)$ . Summing the lengths of all the  $J$ -segments at  $v$ , we count each edge in  $J$  twice and the other edges once; thus, the sum equals  $d_H(v) + d_J(v)$ . Since the lengths of  $J$ -segments are even, so is the sum. Consequently,  $d_H(v)$  and  $d_J(v)$  have the same parity as claimed.  $\square$

The most important reason for us to consider balanced joins, however, is a close relation to perfect matchings:

**Lemma 7.** *Let  $\tilde{G}$  be an  $M$ -contraction of a cubic graph  $G$ , where  $M$  is a perfect matching of  $G$ . A set  $J \subset E(\tilde{G})$  is a balanced join if and only if there is a perfect matching  $M'$  in  $G$  such that*

$$M \cap M' = J.$$

*Proof.* Let  $v_C$  be a vertex of  $\tilde{G}$  and  $J \subset E(\tilde{G})$ . Let  $(h_1, \dots, h_k)$  be a  $J$ -segment at  $v$ . The lemma follows from the observation that  $k$  is even if and only if one can match the vertices of  $G$  incident with  $h_2, \dots, h_{k-1}$  using edges of  $G - M$ .  $\square$

## 4 Balanced joins and the dual graph

There is a relation between balanced joins in an embedded graph  $H$  and independent sets in the dual graph  $H^*$ . Although this relation is not our major concern in the present paper, it is worth a brief investigation which we give in this section.

Recall that if  $H$  is an embedded graph, the dual graph  $H^*$  contains one edge joining faces  $F, F'$  (of the given embedding of  $H$ ) for each edge of  $H$  with the face  $F$  on one side and  $F'$  on the other. In particular, if  $H$  contains an edge with the same face on both sides, then  $H^*$  contains a loop.

We will call a set of faces of an  $H$  *independent* if it corresponds to an independent set in  $H^*$ . Some balanced joins can be obtained from independent sets of faces:

**Proposition 8.** *Let  $I$  be an independent set of faces in an  $M$ -contraction  $H$  of a cubic graph  $G$ . The edges of  $H$  not incident with any face in  $I$  form a balanced join in  $H$ .*

*Proof.* Let  $J$  be set of edges of  $H$  not incident with any face in  $I$ . We need to show that every  $J$ -segment has even length. Consider first a  $J$ -segment

$$[h_1, h_k] = \{h_1, \dots, h_k\} \quad (1)$$

at a vertex  $v$ . Let  $F_i$  be the face of  $H$  delimited by  $h_i$  and  $h_{i+1}$  at  $v$ , where  $i = 1, \dots, k-1$ . Since  $h_1$  is not incident with a face in  $I$ ,  $F_1 \notin I$ . Now  $F_2 \in I$ , because  $h_2 \notin J$ . By the assumption that  $I$  is independent,  $F_3 \notin I$ . Continuing in this way, we conclude that  $F_i \in I$  if and only if  $i$  is even. Since  $F_{k-1} \notin I$ ,  $k$  is even as claimed.

It remains to consider a segment consisting of all the half-edges incident with a vertex  $v$ . As above, we observe that the faces in  $I$  and not in  $I$  must alternate as we traverse around  $v$ , so there must be an even number of them. Consequently,  $|E(v)|$  is even and the proof is complete.  $\square$

It is important to note that not all balanced joins in an embedded graph  $H$  correspond to independent sets in  $H^*$ . In our example from Figure 2, the embedding of  $\tilde{\mathcal{P}}$  shown in Figure 3 has a single face. Thus, the dual graph consists of a single vertex with 5 loops, and the only independent set in  $\tilde{\mathcal{P}}^*$  is  $\emptyset$ . This corresponds to the balanced join  $E(\tilde{\mathcal{P}})$ . However,  $\tilde{\mathcal{P}}$  also contains 5 balanced joins of size 1.

Given Proposition 8, it is not surprising that non-intersecting balanced joins are related to face colorings:

**Proposition 9.** *Let  $\tilde{G}$  be an  $M$ -contraction of a cubic graph  $G$ . If the faces of  $\tilde{G}$  can be properly colored in  $k$  colors, then  $G$  has  $k$  perfect matchings with empty intersection.*

*Proof.* Consider a coloring of the faces in  $k$  colors with color classes  $C_1, \dots, C_k$ . Each  $C_i$  is an independent set of faces. Let  $M'_i$  be the set of edges not incident with any face in  $C_i$ . By Proposition 8, there is a perfect matching  $M_i$  in  $G$  such that  $M_i \cap M = M'_i$ . Since  $C_k$  is independent, each edge of  $M$  is incident with at least one of the sets  $C_i$  with  $i \leq k-1$ . Consequently,

$$M_1 \cap \dots \cap M_{k-1} \subseteq \overline{M}.$$

It follows that the perfect matchings  $M, M_1, \dots, M_{k-1}$  have empty intersection.  $\square$

For  $k = 3$ , one can actually get a stronger result:

**Theorem 10.** *Let  $G$  be a cubic graph with a perfect matching  $M$ . If the faces of some  $M$ -contraction of  $G$  can be properly colored with 3 colors, then  $G$  is 3-edge-colorable.*

The proof is similar to that of Proposition 9 and is omitted. Alternatively, at least under the assumption that every face of an  $M$ -contraction  $\tilde{G}$  is bounded by a circuit, Theorem 10 follows from standard facts on nowhere-zero flows and cycle covers (see, e.g., [9]).

As a curiosity, let us mention the following (probably known) consequence of Theorem 10 and the Grötzsch theorem [5]:

**Proposition 11.** *Let  $G$  be a cyclically 4-edge-connected cubic plane graph with a perfect matching  $M$  such that each circuit of  $\overline{M}$  bounds a face. Then  $G$  is 3-edge-colorable.*

*Proof.* Consider an  $M$ -contraction  $\tilde{G}$  of  $G$ . Since  $\tilde{G}$  is 4-edge-connected, its dual  $\tilde{G}^*$  is triangle-free and hence 3-colorable by the Grötzsch theorem. By Theorem 10,  $G$  is 3-edge-colorable.  $\square$

## 5 A splitting lemma

A basic tool we use in the proof of Theorem 4 is splitting. Let  $G$  be a graph and  $h$  a half-edge incident with a vertex  $v \in V(G)$ . For convenience, we define the *far endvertex* of  $h$  to be the vertex incident with the half-edge associated to  $h$ . Let  $h'$  be another half-edge incident with  $v$ . Furthermore, let  $w$  and  $w'$  be the far endvertices of  $h$  and  $h'$ , respectively.

*Splitting off  $h$  and  $h'$*  is the operation of removing the edges containing  $h$  and  $h'$ , and adding an edge between  $w$  and  $w'$  (a loop if  $w = w'$ ). The resulting graph is denoted by  $G(v; h, h')$ .

The following statement is an equivalent form of the well-known splitting lemma of Fleischner [3] (see also [9, Theorem A.5.2]):

**Lemma 12.** *Let  $v$  be a vertex of a 2-edge-connected graph  $G$  with  $d_G(v) \geq 4$ . Let  $h_0, h_1, h_2 \in E(v)$ . If neither  $G(v; h_0, h_1)$  nor  $G(v; h_0, h_2)$  is 2-edge-connected, then  $\{h_0, h_1, h_2\}$  is a 3-cut in  $G$ .*

Our purpose requires a slightly different kind of a splitting lemma which we derive next. Let  $\mathcal{G}_2$  be the class of all bridgeless graphs with exactly two vertices of odd degree.

**Lemma 13.** *Let  $G \in \mathcal{G}_2$  and let  $h_0, \dots, h_4$  be distinct half-edges incident with a vertex  $z$  of even degree. Let  $w_i$  ( $i = 0, \dots, 4$ ) denote the far endvertex of  $h_i$  and assume that  $w_0$  has odd degree. If neither  $G(z; h_1, h_2)$  nor  $G(z; h_3, h_4)$  is bridgeless, then  $\{h_0, h_1, h_2\}$  or  $\{h_0, h_3, h_4\}$  is a 3-cut in  $G$ .*

*Proof.* Suppose that the statement is false. Choose a bridge  $b'$  in the graph  $G' := G(z; h_1, h_2)$  and a bridge  $b''$  in  $G'' := G(z; h_3, h_4)$ . Since  $G \in \mathcal{G}_2$ , each component of  $G' - b'$  and  $G'' - b''$  contains exactly one vertex of odd degree. We let these vertices be denoted by  $y_1$  and  $y_2$ , where  $y_1 = w_0$ . For  $i \in \{1, 2\}$ , we write  $C'_i$  ( $C''_i$ ) for the vertex set of the component of  $G' - b'$  ( $G'' - b''$ , respectively) that contains  $y_i$ . Finally, we set

$$C_{ij} = C'_i \cap C''_j \quad \text{and} \quad c_{ij} = |\partial_G(C_{ij})|,$$

where  $i, j \in \{1, 2\}$ . Thus,  $y_1 \in C_{11}$  and  $y_2 \in C_{22}$ .

Observe that

$$c_{11} + c_{22} \leq |\partial_G(C'_1)| + |\partial_G(C''_1)| = 6, \quad (2)$$

since each edge contributing to the left hand side is contained in  $\partial_G(C'_1) \cup \partial_G(C''_1)$ , and the edges with a contribution of two (namely, those joining  $C_{11}$  to  $C_{22}$ ) also contribute two to the middle expression.

We distinguish several cases according to the location of the vertex  $z$ . Assume first that  $z \in C_{11}$ . Thus,  $\{h_1, \dots, h_4\} \subset \partial_G(C_{11})$ . Since  $C_{11}$  contains exactly one vertex of odd degree,  $c_{11}$  is odd and hence  $c_{11} \geq 5$ . By (2),  $c_{22} \leq 1$  and  $G$  contains a bridge, which is a contradiction. In a similar way, we exclude the possibility that  $z \in C_{22}$ .

By symmetry, we may therefore assume that  $z \in C_{21}$ . Let  $e_i$  ( $i = 0, 1, 2$ ) be the edge containing  $h_i$ . Since  $w_0 = y_1 \in C_{11}$ , we have

$$e_0 \in \partial_G(C'_1) = \{e_1, e_2, b'\},$$

whence  $e_0 = b'$ . In a bridgeless graph,  $|\partial_G(X)| = 3$  implies that  $\partial_G(X)$  is a 3-cut. We conclude that  $\{e_0, e_1, e_2\}$  is a 3-cut in  $G$ , proving the assertion.  $\square$

## 6 The 5-line conjecture for graphs of oddness two

Let us begin with a notion that can be used to reformulate Conjecture 3. We will say that a set  $X \subset E(G)$  is *sparse* if  $X$  contains no odd cut.

**Observation 14.** *A set  $X \subset E(G)$  is sparse if and only if there exists a join  $J$  with  $X \cap J = \emptyset$ .*

In this section, we prove that every embedded graph from the class  $\mathcal{G}_2$  admits a sparse balanced join. At the end of the section, we derive Theorem 4 as a corollary.

Let  $G$  be a graph,  $e \in E(G)$  and let  $z$  be an endvertex of  $e$ . An unordered pair  $\{f, f'\}$  of edges is *near  $e$  at  $z$*  if

- (1)  $e \notin \{f, f'\}$ ,

- (2)  $f$  and  $f'$  are consecutive at  $z$  and one of them immediately follows or precedes  $e$  at  $z$ .

In the above definition, we allow  $f = f'$  if  $f$  is a loop whose ends are consecutive at  $z$ .

**Lemma 15.** *Let  $e$  be an edge of a graph  $G$  with endvertex  $z$ . Let  $G'$  be a bridgeless graph obtained by splitting off a pair of edges near  $e$  at  $z$ . If  $G'$  has a sparse balanced join containing  $e$ , then so does  $G$ .*

*Proof.* Assume we split off the edges  $f$  and  $h$  following  $e$  at  $z$  (in this order) to obtain  $G'$ . Let  $J'$  be a sparse balanced join in  $G'$  with  $e \in J'$ , and let  $J$  be the corresponding join in  $G$ . Note that  $e \in J$ . It is easy to compare the sizes of the  $J'$ -segments at  $z$  in  $G'$  and the  $J$ -segments in  $G$ . They are the same, except that:

- if  $f, h \notin J$ , then the  $J$ -segment at  $z$  in  $G$  containing  $f$  and  $h$  is larger by 2 than the corresponding  $J'$ -segment in  $G'$ ,
- otherwise, there are two extra  $J$ -segments at  $z$  in  $G$  whose size is 2 (delimited by  $e, f$  and  $f, h$ ).

In either case, the lengths of all the  $J$ -segments are even, which means that  $J$  is balanced.

Assume that  $J$  contains an odd cut  $C$ . Necessarily,  $\{f, h\} \subset J$ . Let  $C' = C - \{f, h\}$ . Since  $C'$  is an odd cut in  $G'$ , we have  $C' \not\subset J'$  and hence  $C \not\subset J$ , a contradiction. It follows that  $J$  is sparse. The proof is finished.  $\square$

**Theorem 16.** *Let  $G \in \mathcal{G}_2$  be an embedded graph and  $e \in E(G)$  an edge incident with an odd degree vertex. Then  $G$  admits a sparse balanced join  $J$  such that  $e \in E(J)$ .*

*Proof.* Let  $G$  be a counterexample with as few edges as possible. Since  $G$  is bridgeless and  $G \in \mathcal{G}_2$ , it has at least 3 edges. If  $G$  has exactly 3 edges, it consists of two vertices joined by three parallel edges, and the sparse balanced join  $\{e\}$  provides a contradiction.

We may thus assume that  $|E(G)| > 3$  and that the assertion holds for graphs with fewer edges. We may also assume that the minimum degree in  $G$  is at least 3 and that  $e$  has only one endvertex ( $u$ , say) of odd degree. Let  $z$  denote the other endvertex of  $e$ , and  $v$  denote the remaining vertex of odd degree in  $G$ .

By Lemma 15 and the minimality of  $G$ , it is not possible to split off any pair of half-edges near  $e$  at  $z$  so as to obtain a bridgeless graph. Enumerate the half-edges incident with  $z$ , in the rotation order, as  $h_0, \dots, h_{k-1}$ , where  $h_0$  is a part of  $e$ . By the above, the degree  $k = d_G(z)$  of  $z$  is even and not equal to 2.

Assume now that  $k = 4$ . By Lemma 12, either  $G(z; h_1, h_2)$  or  $G(z; h_2, h_3)$  is a bridgeless graph. Since both  $\{h_1, h_2\}$  and  $\{h_2, h_3\}$  are near  $e$  at  $z$ , we get a contradiction. Thus,  $k \geq 6$ .

Lemma 13 implies that at least one of the sets  $\{h_0, h_1, h_2\}$  or  $\{h_0, h_{k-1}, h_{k-2}\}$  is a 3-cut. By symmetry, we may assume that  $\{h_0, h_1, h_2\}$  is a 3-cut. Since  $d_G(z) \geq 6$ ,  $z$  is a cutvertex. Moreover, there are exactly three edges joining  $z$  to the component  $X$  of  $G - z$  containing  $u$ . Since  $|\partial_G(V(X))|$  is odd,  $v$  (the other vertex of odd degree in  $G$ ) is not contained in  $X$ .

Set  $G' = G - V(X)$ . Note that the degree of  $z$  in  $G'$  is odd (namely,  $k - 3$ ) and that  $G' \in \mathcal{G}_2$ . By the minimality of  $G$ , there is a balanced join  $J'$  in  $G'$  such that  $h_3 \in J'$  and  $J'$  contains no odd cut in  $G'$ . Clearly,

$$J = J' \cup \{e\}$$

is a balanced join in  $G$ . Furthermore, since every odd cut in  $G$  is contained either in  $G - V(X)$  or in  $G[V(X) \cup \{z\}]$ ,  $J$  is sparse. Thus,  $J$  has the properties stated in the theorem, a contradiction.  $\square$

We can now derive Theorem 4 as a corollary of Theorem 16.

*Proof of Theorem 4.* Let  $G$  be a cubic bridgeless graph of oddness two. Let  $M_1$  be a perfect matching such that the 2-factor  $\overline{M_1}$  has exactly two odd components. Choose an orientation  $o$  of  $\overline{M_1}$  and consider the  $M_1$ -contraction  $H = G_{M_1, o}$ . Note that  $H \in \mathcal{G}_2$ . By Theorem 16,  $H$  admits a sparse balanced join  $J_2$ .

Observation 7 implies that  $J_2$  can be extended to a perfect matching  $M_2$  in  $G$  such that  $M_1 \cap M_2 = J_2$ . Since  $J_2$  is sparse in  $H$ ,  $M_1 \cap M_2$  is sparse in  $G$ . By Observation 14, there is a join  $J$  in  $G$  such that  $M_1 \cap M_2 \cap J = \emptyset$ . This completes the proof.  $\square$

## 7 Concluding remarks

There are a number of natural questions about balanced joins. The first of them concerns a possible extension of Theorem 16 to embedded graphs with more than two odd degree vertices. The statement of the theorem does not hold for all bridgeless graphs: for instance, it is easy to see that the Petersen graph  $\mathcal{P}$ , together with an arbitrary rotation system, has no sparse join. However, the following may be true:

**Conjecture 17.** *Let  $H$  be an embedded graph with  $\lambda^{\text{odd}}(H) \geq 5$ . Then  $H$  contains a sparse balanced join.*

A natural approach to the Fan–Raspau conjecture (Conjecture 1) along the lines of our proof of Theorem 4 would require finding two disjoint balanced joins in an embedded graph. Again, the Petersen graph  $\mathcal{P}$  does not even have two disjoint joins, but we find it plausible that higher odd edge-connectivity helps.

**Problem 18.** *Is there an integer  $k$  such that every embedded graph  $H$  of odd edge-connectivity  $\lambda^{\text{odd}}(H) \geq k$  contains two disjoint balanced joins?*

Observe that if the balanced joins in Problem 18 exist, they are necessarily sparse.

We may go one step further and ask about partitions into balanced joins. It is interesting to consider ‘the’  $M$ -contraction  $\tilde{\mathcal{P}}$  of the Petersen graph in Figure 2b. Since  $E(\tilde{\mathcal{P}})$  and any set consisting of a single edge are balanced joins, the edge set of  $\tilde{\mathcal{P}}$  can be partitioned into 1 or 5 balanced joins. However, there is no partition of  $E(\tilde{\mathcal{P}})$  into 3 balanced joins.

**Problem 19.** *Is there a function  $f$  such that for every odd  $\ell$ , the edge set of each embedded graph  $H$  with  $\lambda^{\text{odd}}(H) \geq f(\ell)$  can be partitioned into  $\ell$  balanced joins?*

## References

- [1] R. Diestel, *Graph Theory*, Springer-Verlag, New York, 2000.
- [2] G. Fan and A. Raspaud, Fulkerson’s Conjecture and circuit covers, *J. Combin. Theory Ser. B* **61** (1994), 133–138.
- [3] H. Fleischner, Eine gemeinsame Basis für die Theorie der eulerschen Graphen und den Satz von Petersen, *Monatsh. Math.* **81** (1976), 267–278.
- [4] D. R. Fulkerson, Blocking and anti-blocking pairs of polyhedra, *Math. Programming* **1** (1971), 168–194.
- [5] H. Grötzsch, Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel, *Wiss. Z. Martin-Luther-Univ. Halle–Wittenberg Math.-Natur. Reihe* **8** (1959), 109–120.
- [6] F. Holroyd and M. Škoviera, Colouring of cubic graphs by Steiner triple systems, *J. Combin. Theory Ser. B* **91** (2004), 57–66.
- [7] E. Máčajová and M. Škoviera, Fano colourings of cubic graphs and the Fulkerson conjecture, *Theor. Comput. Sci.* **349** (2005), 112–120.
- [8] B. Mohar and C. Thomassen, *Graphs on Surfaces*, Johns Hopkins University Press, Baltimore, MD, 2001.
- [9] C.-Q. Zhang, *Integer Flows and Cycle Covers of Graphs*, M. Dekker, New York, 1997.