

# Planar 4-critical graphs with four triangles

Oleg V. Borodin <sup>\*</sup>      Zdeněk Dvořák <sup>†</sup>  
Alexandr V. Kostochka <sup>‡</sup>      Bernard Lidický <sup>§</sup>  
Matthew Yancey <sup>¶</sup>

July 20, 2013

## Abstract

By the Grünbaum-Aksenov Theorem (extending Grötzsch's Theorem) every planar graph with at most three triangles is 3-colorable. However, there are infinitely many planar 4-critical graphs with exactly four triangles. We describe all such graphs. This answers a question of Erdős from 1990.

---

<sup>\*</sup>Sobolev Institute of Mathematics and Novosibirsk State University, Novosibirsk 630090, Russia. E-mail: [brdnoleg@math.nsc.ru](mailto:brdnoleg@math.nsc.ru). Research of this author is supported in part by grants 12-01-0044. and 12-01-00631 of the Russian Foundation for Basic Research.

<sup>†</sup>Computer Science Institute of Charles University, Prague, Czech Republic. E-mail: [rakdver@iuuk.mff.cuni.cz](mailto:rakdver@iuuk.mff.cuni.cz). Supported the Center of Excellence – Inst. for Theor. Comp. Sci., Prague (project P202/12/G061 of Czech Science Foundation), and by project LH12095 (New combinatorial algorithms - decompositions, parameterization, efficient solutions) of Czech Ministry of Education.

<sup>‡</sup>University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA and Sobolev Institute of Mathematics, Novosibirsk 630090, Russia. E-mail: [kostochk@math.uiuc.edu](mailto:kostochk@math.uiuc.edu). Research of this author is supported in part by NSF grant DMS-0965587.

<sup>§</sup>University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. E-mail: [lidicky@illinois.edu](mailto:lidicky@illinois.edu)

<sup>¶</sup>University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. E-mail: [yancey1@illinois.edu](mailto:yancey1@illinois.edu). Research of this author is supported by National Science Foundation grant DMS 08-38434 “EMSW21-MCTP: Research Experience for Graduate Students.”




Figure 1: A quasi-edge  $H_0$  and  $\text{Pl}_4$ -graphs  $H_1$  and  $H_2$

## 1 Introduction

The classical Grötzsch's Theorem [14] says that every planar triangle-free graph is 3-colorable. The following refinement of it is known as the Grünbaum-Aksenov Theorem (the original proof of Grünbaum [15] was incorrect, and Aksenov [1] fixed the proof).

**Theorem 1** ([1, 7, 15]). *Let  $G$  be a planar graph containing at most three triangles. Then  $G$  is 3-colorable.*

The example of the complete 4-vertex graph  $K_4$  shows that “three” in Theorem 1 cannot be replaced by “four”. But maybe there are not many plane 4-critical graphs with exactly four triangles ( $\text{Pl}_4$ -graphs, for short)?

It turned out that there are many. Havel [16] presented a  $\text{Pl}_4$ -graph  $H_1$  (see Figure 1) in which the four triangles had no common vertices. He used the quasi-edge  $H_0 = H_0(u, v)$  (on the left of Figure 1), that is, a graph in each 3-coloring of which the vertices  $u$  and  $v$  must have distinct colors. The graph  $H_1$  is obtained from  $K_4$  by replacing the edges  $v_1u_1$  and  $v_2u_2$  with copies of the quasi-edge  $H_0$ . Then Sachs [18] in 1972 asked whether it is true that in every non-3-colorable planar graph  $G$  with exactly four triangles and no separating triangles these triangles can be partitioned into two pairs so that in each pair the distance between the triangles is less than two.

Aksenov and Mel'nikov [5, 6] answered the question in the negative by constructing a  $\text{Pl}_4$ -graph  $H_2$  (see Figure 1) in which the distance between any two of the four triangles is at least two. Moreover, they constructed two infinite series of  $\text{Pl}_4$ -graphs. Aksenov [2] was studying  $\text{Pl}_4$ -graphs in the seventies. According to Steinberg [19], Erdős in 1990 asked for a description




Figure 2: Some members of  $\mathcal{TW}$ .

of  $\text{Pl}_4$ -graphs again. Borodin [7] remarks that he knows 15 infinite families of  $\text{Pl}_4$ -graphs. In his survey [8], he mentions the problem of describing  $\text{Pl}_4$ -graphs among unsolved problems on 3-coloring of plane graphs.

First, we give a description of the  $\text{Pl}_4$ -graphs without 4-faces, which we call  $\text{Pl}_{4,4f}$ -graphs. Thomas and Walls [20] constructed an infinite family  $\mathcal{TW}$  of  $\text{Pl}_{4,4f}$ -graphs; the first five graphs in  $\mathcal{TW}$  are depicted in Figure 2 (note that  $T_2$  and  $T_3$  are isomorphic graphs, but their drawings are different). If an edge  $e$  of a graph belongs to exactly two triangles, we say that  $e$  is a *diamond* edge. Each graph in  $\mathcal{TW}$  contains two disjoint diamond edges, drawn in bold in Figure 2. We define the class  $\mathcal{TW}$  in terms of Ore-compositions.

An *Ore-composition*  $O(G_1, G_2)$  of graphs  $G_1$  and  $G_2$  is a graph obtained as follows: delete some edge  $xy$  from  $G_1$ , split some vertex  $z$  of  $G_2$  into two vertices  $z_1$  and  $z_2$  of positive degree, and identify  $x$  with  $z_1$  and  $y$  with  $z_2$ . If  $xy$  is a diamond edge of  $G_1$  and  $G_2$  is  $K_4$ , then we say that  $O(G_1, G_2)$  is a *diamond expansion* of  $G_1$ . The class  $\mathcal{TW}$  consists of all graphs that can be obtained from  $K_4$  by diamond expansions.

Note that  $H_1$  is a  $\text{Pl}_{4,4f}$ -graph but is not in  $\mathcal{TW}$ . A graph is  $k$ -Ore if it is obtained from a set of copies of  $K_k$  by a sequence of Ore-compositions. It was proved in [17] that every  $k$ -Ore graph is  $k$ -critical. A partial case of Theorem 6 in [17] is the following.

**Theorem 2** ([17]). *Let  $G$  be an  $n$ -vertex 4-critical graph. Then  $|E(G)| \geq \frac{5n-2}{3}$ . Moreover,  $|E(G)| = \frac{5n-2}{3}$  if and only if  $G$  is a 4-Ore graph.*

We will see below that every  $\text{Pl}_{4,4f}$ -graph is a 4-Ore graph. On the other hand, our first result says the following.

**Theorem 3.** *Every 4-Ore graph has at least four triangles. Moreover, a 4-Ore graph  $G$  has exactly four triangles if and only if  $G$  is a  $\text{Pl}_{4,4f}$ -graph.*




Figure 3: Some members of  $\mathcal{TW}_1$ .

This reduces a “topological” result on plane graphs to a result on abstract graphs. And it turns out that  $\text{Pl}_{4,4f}$ -graphs do not differ much from the Thomas-Walls graphs. Let  $\mathcal{TW}_1$  denote the graphs obtained from a graph in  $\mathcal{TW}$  by replacing a diamond edge by the Havel’s quasi-edge  $H_0$  (see Figure 3). Let  $\mathcal{TW}_2$  denote the graphs obtained from a graph in  $\mathcal{TW}_1$  by replacing a diamond edge by the Havel’s quasi-edge  $H_0$  (see Figure 4). Note that  $H_0$  contains no diamond edges, and thus each graph in  $\mathcal{TW}_2$  can be obtained from a graph in  $\mathcal{TW}$  by replacing two vertex-disjoint diamond edges by the Havel’s quasi-edge  $H_0$ .

**Theorem 4.** *The class of  $\text{Pl}_{4,4f}$ -graphs is equal to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ .*

The  $\text{Pl}_4$ -graphs may have arbitrarily many 4-faces. Let  $G$  be a plane graph, let  $C_P = xz'yx'zy'$  be a 6-cycle in  $G$  and let  $\Delta$  be the closed disk bounded by  $C_P$ . Let  $P$  be the subgraph of  $G$  consisting of the vertices and edges drawn in  $\Delta$ . If all neighbors of the vertices  $x'$ ,  $y'$  and  $z'$  belong to  $P$  and all faces of  $G$  contained in  $\Delta$  have length 4, then we say that  $P$  is a *patch*. The patch  $P$  is *critical* if  $x'$ ,  $y'$  and  $z'$  have degree at least 3 and every




Figure 4: Some members of  $\mathcal{TW}_2$ .

4-cycle in  $P$  bounds a face. We will see that if  $G_0$  is a  $\text{Pl}_4$ -graph and a vertex  $v \in V(G_0)$  has exactly 3 neighbors  $x, y$  and  $z$ , then the graph  $G_v$  obtained from  $G_0 - v$  by inserting a critical patch  $P$  with boundary  $C_P = xz'yx'zy'$  (where  $x', y'$  and  $z'$  are new vertices) is again a  $\text{Pl}_4$ -graph. For example, the graph  $H_2$  in Figure 1 is obtained from the graph  $H_1$  by replacing the vertices  $r$  and  $s$  with patches bounded by the cycles  $r_1 \dots r_6$  and  $s_1 \dots s_6$ , respectively. This gives a way to construct from every  $\text{Pl}_{4,4f}$ -graph an infinite family of  $\text{Pl}_4$ -graphs. Our main result is that every  $\text{Pl}_4$ -graph can be obtained this way.

**Theorem 5.** *A plane 4-critical graph has exactly four triangles if and only if it is obtained from a  $\text{Pl}_{4,4f}$ -graph by replacing several (possibly zero) non-adjacent 3-vertices with critical patches.*

Thus, even though there are infinitely many  $\text{Pl}_4$ -graphs, we know the structure of all of them. This fully answers Erdős' question from 1990. In particular, the result yields that Sachs had the right intuition in 1972: his question has positive answer if we replace “less than two” with “at most two.” Also, Aksenov and Mel'nikov [6] conjectured, in particular, that  $H_1$  is the unique smallest  $\text{Pl}_4$ -graph with the minimum distance 1 between triangles and  $H_2$  is the unique smallest  $\text{Pl}_4$ -graph with the minimum distance 2 between triangles. Our description confirms this.

Havel [16] asked the following question: Does there exist a constant  $C$  such that every planar graph with the minimal distance between triangles at least  $C$  is 3-colorable? The graph  $H_2$  shows that  $C \geq 3$ , and a further example of Aksenov and Mel'nikov [6] shows that  $C \geq 4$ , which is the best known lower bound. The existence of (large)  $C$  was recently proved by Dvořák, Král' and Thomas [9]. It is conjectured by Borodin and Raspaud that  $C = 4$  is sufficient, and our result confirms this conjecture for graphs with four triangles. For more details, see a recent survey of Borodin [8].

The proof of Theorem 5 can be converted to a polynomial-time algorithm to find a 3-coloring of a planar graph with four triangles or to decide that no such coloring exists. However, a more general algorithm of Dvořák, Král' and Thomas [10] can also be used, and thus we do not provide further details.

The structure of the paper is as follows. In the next section we study the structure of 4-Ore graphs. In Section 3 we prove Theorems 3 and 4. In the last section, we describe all  $\text{Pl}_4$ -graphs by proving Theorem 5.

## 2 The structure of 4-Ore graphs

In this section, we study 4-Ore graphs with few triangles. The following claim is simply a reformulation of the definition of a  $k$ -Ore graph.

**Claim 6.** *Every  $k$ -Ore graph  $G \neq K_k$  has a separating set  $\{x, y\}$  and two vertex subsets  $A$  and  $B$  such that*

- $A \cap B = \{x, y\}$ ,  $A \cup B = V(G)$  and no edge of  $G$  connects  $A \setminus \{x, y\}$  with  $B \setminus \{x, y\}$ ,
- $x$  and  $y$  are non-adjacent in  $G$  and have no common neighbor in  $B$ ,
- the graph  $G'$  obtained from  $G[A]$  by adding the edge  $xy$  is a  $k$ -Ore graph, and
- the graph  $G''$  obtained from  $G[B]$  by identifying  $x$  with  $y$  into a new vertex  $x * y$  is a  $k$ -Ore graph.

Our first goal is to obtain a similar decomposition when we restrict ourselves to 4-Ore graphs with 4 triangles (Theorem 11 below).

**Claim 7.** *Every edge in each 4-Ore graph is contained in at most 2 triangles.*

*Proof.* We prove the claim by induction on the order of a graph. Let  $G$  be a 4-Ore graph and  $uv$  its edge and assume that the claim holds for all graphs with less than  $|V(G)|$  vertices.

Each edge of  $K_4$  is contained in exactly two triangles, and thus we can assume that  $G \neq K_4$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Claim 6. Since  $u$  is adjacent to  $v$ , either  $\{u, v\} \subset A$  or  $\{u, v\} \subset B$ . Let  $G_0 \in \{G[A], G[B]\}$  be the graph containing the edge  $uv$ , and let  $G'_0 \in \{G', G''\}$  be the corresponding 4-Ore graph. Let  $u'v'$  be the edge of  $G'_0$  corresponding to  $uv$ . Every triangle of  $G$  containing  $uv$  maps to a triangle in  $G'_0$  containing  $u'v'$ . Since  $|V(G'_0)| < |V(G)|$ , the edge  $u'v'$  is contained in at most two triangles in  $G'_0$  by induction, and thus  $uv$  is contained in at most two triangles in  $G$ .  $\square$

For a graph  $G$ , let  $t(G)$  denote the number of triangles in  $G$ .

**Claim 8.** *If  $G$  is a 4-Ore graph, then  $t(G) \geq 4$ . For every vertex  $z \in V(G)$ , every graph  $G_z$  obtained from  $G$  by splitting  $z$  satisfies  $t(G_z) \geq 2$ . Furthermore, if  $G \neq K_4$ , then  $t(G-z) \geq 2$ , and if  $G = K_4$ , then  $t(G-z) = 1$ .*

*Proof.* Let  $G$  be a 4-Ore graph. We proceed by induction and assume that the claim holds for all graphs with less than  $|V(G)|$  vertices. Since the claim holds for  $K_4$ , we can assume that  $G \neq K_4$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Claim 6. By induction hypothesis and Claim 7,  $G[A] = G' - xy$  has at least  $t(G') - 2 \geq 2$  triangles. Furthermore,  $G[B]$  is obtained from  $G''$  by splitting a vertex, and thus it has at least two triangles by induction. It follows that  $G$  has at least four triangles.

Consider now a vertex  $z$  of  $G$ . If  $z \notin \{x, y\}$ , then  $G - z$  contains  $G[A]$  or  $G[B]$  as a subgraph, and thus  $t(G - z) \geq 2$ . If  $z \in \{x, y\}$ , say  $z = x$ , then  $G - z$  contains  $G' - x$  and  $G'' - x * y$  as vertex-disjoint subgraphs, and by induction each of them has a triangle; hence,  $t(G - z) \geq 2$ . Finally, any graph  $G_z$  obtained from  $G$  by splitting  $z$  contains  $G - z$  as a subgraph, and thus  $t(G_z) \geq t(G - z) \geq 2$ .  $\square$

**Claim 9.** *If  $G$  is a 4-Ore graph with  $t(G) \geq 5$ , then  $t(G - u - v) \geq 1$  for each  $u, v \in V(G)$ .*

*Proof.* We proceed by induction and assume that the claim holds for all graphs with less than  $|V(G)|$  vertices. Note that  $G \neq K_4$ , since  $t(G) \geq 5$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Claim 6. If  $u, v \in A$ , then  $G'' - x * y$  is a subgraph of  $G - u - v$  and  $t(G - u - v) \geq t(G'' - x * y) \geq 1$  by Claim 8. Hence, by symmetry we can assume that  $u \in B \setminus \{x, y\}$ . Suppose that  $v \in B$ . We can assume that  $v \neq x$ , and thus  $G' - y$  is a subgraph of  $G - u - v$ . Again, Claim 8 implies that  $t(G - u - v) \geq t(G' - y) \geq 1$ .

Finally, consider the case that  $u \in B \setminus \{x, y\}$  and  $v \in A \setminus \{x, y\}$ . Since  $t(G) \geq 5$ , either  $G[A]$  or  $G[B]$  contains at least three triangles. In the former case, let  $G_0 = G[A]$ ,  $G'_0 = G'$ ,  $r = v$  and  $s = x$ . In the latter case, let  $G_0 = G[B]$ ,  $G'_0 = G''$ ,  $r = u$  and  $s = x * y$ . If  $r$  is contained in at most two triangles in  $G_0$ , then  $t(G - u - v) \geq t(G_0 - r) \geq t(G_0) - 2 \geq 1$ ; hence, assume that  $r$  is contained in at least 3 triangles in  $G_0$ , and thus also in  $G'_0$ . Note that  $G'_0 \neq K_4$ , since  $t(G_0) \geq 3$ . By Claim 8, we have  $t(G'_0) \geq t(G'_0 - r) + 3 \geq 5$ . Note that  $G'_0 - r - s$  is a subgraph of  $G - u - v$ , and thus  $t(G - u - v) \geq t(G'_0 - r - s) \geq 1$  by induction.  $\square$

**Claim 10.** *If  $G$  is a 4-Ore graph with  $t(G) \geq 5$ , then  $t(G - v) \geq 3$  for each  $v \in V(G)$ .*

*Proof.* Note that  $G \neq K_4$ , since  $t(G) \geq 5$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Claim 6. Since  $t(G) \geq 5$ , either  $G[A]$  or  $G[B]$  contains at least three triangles. In the former case, let  $G_0 = G[A]$ ,  $G_1 = G[B]$ ,  $G'_0 = G'$  and  $s = x$ . In the latter case, let  $G_0 = G[B]$ ,  $G_1 = G[A]$ ,  $G'_0 = G''$  and  $s = x * y$ . Since  $G_0$  has at least three triangles, it follows that  $G'_0 \neq K_4$ . If  $v \notin V(G_0)$ , then  $t(G - v) \geq t(G_0) \geq 3$ . Therefore, assume that  $v \in V(G_0)$ . If  $v \in \{x, y\}$ , then  $G' - v$  and  $G'' - x * y$  are vertex-disjoint subgraphs of  $G - v$ , and since at least one of  $G'$  and  $G''$  is not equal to  $K_4$ , we have  $t(G - v) \geq t(G' - v) + t(G'' - x * y) \geq 3$  by Claim 8.

Finally, suppose that  $v \notin \{x, y\}$ , and thus  $G_1$  is a subgraph of  $G - v$ . By Claim 8, there are at least two triangles in  $G_1$ . We claim that  $G_0 - v$  contains a triangle. This is clear if  $v$  belongs to at most two triangles in  $G_0$ , since  $t(G_0) \geq 3$ . Otherwise,  $v$  belongs to at least three triangles in  $G_0$ , and thus also in  $G'_0$ . By Claim 8, we have  $t(G'_0) \geq t(G'_0 - v) + 3 \geq 5$ . We conclude that  $t(G_0 - v) \geq t(G'_0 - v - s) \geq 1$  by Claim 9. Therefore,  $t(G - v) = t(G_0 - v) + t(G_1) \geq 3$ .  $\square$

A *4,4-Ore graph* is a 4-Ore graph with exactly 4 triangles. The main result of this section is the following.

**Theorem 11.** *Suppose  $G$  is a 4,4-Ore graph distinct from  $K_4$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Claim 6. Then both  $G'$  and  $G''$  are 4,4-Ore graphs,  $xy$  is a diamond edge of  $G'$ , and  $t(G[B]) = 2$ . Furthermore, if  $G'' \neq K_4$ , then  $x * y$  belongs to exactly two triangles of  $G''$ .*

*Proof.* Note that  $t(G[A]) \geq 2$  and  $t(G[B]) \geq 2$  by Claims 7 and 8, and since  $t(G[A]) + t(G[B]) = t(G) = 4$ , it follows that  $t(G[A]) = t(G[B]) = 2$ . If  $t(G') \geq 5$ , then we would have  $t(G[A]) \geq t(G' - x) \geq 3$  by Claim 10. If  $t(G'') \geq 5$ , then we would have  $t(G[B]) \geq t(G'' - x * y) \geq 3$  by Claim 10. It follows that  $t(G') \leq 4$  and  $t(G'') \leq 4$ , and by Claim 8, we conclude that  $t(G') = t(G'') = 4$ , i.e., both  $G'$  and  $G''$  are 4,4-Ore graphs.

Since  $xy$  belongs to  $t(G') - t(G[A]) = 2$  triangles in  $G'$ , it is a diamond edge. Similarly,  $x * y$  belongs to at least  $t(G'') - t(G[B]) = 2$  triangles in  $G''$ . Furthermore, if  $G'' \neq K_4$  and  $x * y$  belonged to at least three triangles, then we would have  $t(G'') \geq t(G'' - x * y) + 3 \geq 5$  by Claim 8, which is a contradiction.  $\square$



### 3 A description of 4, 4-Ore graphs and $\text{Pl}_{4,4f}$ -graphs

With Theorem 11, it is easy to characterize all 4, 4-Ore graphs. The *Moser spindle* is the Ore composition of two  $K_4$ 's, depicted in Figure 2 as  $M$ .

**Lemma 12.** *Every 4, 4-Ore graph belongs to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ .*

*Proof.* Let  $G$  be a 4, 4-Ore graph. We proceed by induction and assume that every 4, 4-Ore graph with less than  $|V(G)|$  vertices belongs to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ . Note that  $K_4 \in \mathcal{TW}$ , and thus we can assume that  $G \neq K_4$ . Let  $\{x, y\}$ ,  $A$ ,  $B$ ,  $G'$  and  $G''$  be as in Theorem 11. By induction hypothesis, we have  $G', G'' \in \mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ . Since  $G'$  has a diamond edge  $xy$  and  $G''$  has a vertex  $x * y$  belonging to at least two triangles, we conclude that  $G', G'' \notin \mathcal{TW}_2$ . Since  $G[B]$  has exactly two triangles, an inspection of the graphs in  $\mathcal{TW} \cup \mathcal{TW}_1$  (see Figures 2 and 3) shows that either

- $G''$  is the Moser spindle,  $x * y$  is its vertex of degree four,  $G[B]$  is the Havel's quasiedge  $H_0$  and  $x$  and  $y$  are its vertices of degree two, or
- $x$  has degree two in  $G[B]$ ,  $y$  has degree one in  $G[B]$  and  $x * y$  is incident with a diamond edge  $(x * y)z$  of  $G''$ , where  $z$  is the neighbor of  $y$  in  $G[B]$ .

In the former case,  $G$  is obtained from  $G_1$  by replacing a diamond edge with the Havel's quasiedge  $H_0$ , and thus  $G \in \mathcal{TW}_1 \cup \mathcal{TW}_2$ . In the latter case, observe that the described Ore-composition of graphs from  $\mathcal{TW} \cup \mathcal{TW}_1$  results in a graph from  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ .  $\square$

We can now describe  $\text{Pl}_{4,4f}$ -graphs.

*Proof of Theorems 4 and 3.* By Claim 8, every 4-Ore graph has at least four triangles. Therefore, it suffices to prove that the following classes are equal to each other:

- 4, 4-Ore graphs,
- $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ , and
- $\text{Pl}_{4,4f}$ -graphs.

By Lemma 12, every 4, 4-Ore graph belongs to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ . Every graph in  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$  is planar, 4-critical (since it is 4-Ore) and has four triangles, and thus it is a  $\text{Pl}_{4,4f}$ -graph. Therefore, it suffices to show that every  $\text{Pl}_{4,4f}$ -graph  $G$  is 4, 4-Ore. Since  $G$  has four triangles, we only need to prove that  $G$  is 4-Ore. Consider a plane drawing of  $G$  without 4-faces. Let  $e$ ,  $n$  and  $s$  be the number of edges, vertices and faces of the drawing of  $G$ , respectively. Note that  $G$  has at most four triangular faces and all other faces of  $G$  have length at least 5. Therefore,  $G$  has at least  $\frac{1}{2}(5(s-4) + 3 \cdot 4)$  edges. Note that  $G$  is connected (since it is 4-critical), and thus  $s = e + 2 - n$  by Euler's formula. It follows that  $e \leq (5n - 2)/3$ , and  $G$  is 4-Ore by Theorem 2.  $\square$

## 4 A description of $\text{Pl}_4$ -graphs

We are going to characterize planar 4-critical graphs with 4 triangles. To deal with short separating cycles, we use the notion of criticality with respect to a subgraph. Let  $G$  be a graph and  $C$  be its (not necessarily induced) proper subgraph. We say that  $G$  is  $C$ -critical (for 3-coloring) if for every proper subgraph  $H \subset G$  such that  $C \subseteq H$ , there exists a 3-coloring of  $C$  that extends to a 3-coloring of  $H$ , but not to a 3-coloring of  $G$ .

Notice that 4-critical graphs are exactly  $C$ -critical graphs with  $C = \emptyset$ . Furthermore, it is easy to see that if  $F$  is a 4-critical graph and  $F = G \cup G'$ , where  $C = G \cap G'$ , then either  $G = C$  or  $G$  is  $C$ -critical. We mainly use the following reformulation.

**Lemma 13** (Dvořák et al. [11]). *Let  $G$  be a plane graph and let  $\Lambda$  be a connected open region of the plane whose boundary is equal to a cycle  $C$  of  $G$ , such that  $\Lambda$  is not a face of  $G$ . Let  $H$  be the subgraph of  $G$  drawn in the closure of  $\Lambda$ . If  $G$  is 4-critical, then  $H$  is  $C$ -critical.*

This is useful in connection with the following result of Gimbel and Thomassen [13], which was also obtained independently by Aksenov et al. [3].

**Theorem 14** (Gimbel and Thomassen [13]). *Let  $G$  be a plane triangle-free graph with the outer face bounded by a cycle  $C$  of length at most 6. If  $G$  is  $C$ -critical, then  $C$  is a 6-cycle and all internal faces of  $G$  have length four.*

Furthermore, they also exactly characterized the colorings of  $C$  that do not extend to  $G$ .




Figure 5: Critical graphs with a precolored 7-face.

**Theorem 15** (Gimbel and Thomassen [13]). *Let  $G$  be a plane graph with the outer face bounded by a cycle  $C = c_1 \dots c_6$  of length 6, such that all other faces of  $G$  have length 4. A 3-coloring  $\varphi$  of  $G[V(C)]$  does not extend to a 3-coloring of  $G$  if and only if  $\varphi(c_1) = \varphi(c_4)$ ,  $\varphi(c_2) = \varphi(c_5)$  and  $\varphi(c_3) = \varphi(c_6)$ .*

An analogous result for a 7-cycle was obtained by Aksenov et al. [4]. Their paper is in Russian. For a shorter proof in English see [12].

**Theorem 16** (Aksenov et al. [4]). *Let  $G$  be a plane triangle-free graph with the outer face bounded by a cycle  $C = c_1 \dots c_7$  of length 7. The graph  $G$  is  $C$ -critical and  $\varphi : V(C) \rightarrow \{1, 2, 3\}$  is a 3-coloring of  $C$  that does not extend to a 3-coloring of  $G$  if and only if  $G$  contains no separating cycles of length at most five and one of the following propositions is satisfied up to relabelling of vertices (see Figure 5 for an illustration).*

- (a) *The graph  $G$  consists of  $C$  and the edge  $c_1c_5$ , and  $\varphi(c_1) = \varphi(c_5)$ .*
- (b) *The graph  $G$  contains a vertex  $v$  adjacent to  $c_1$  and  $c_4$ , the cycle  $c_1c_2c_3c_4v$  bounds a 5-face and every face drawn inside the 6-cycle  $vc_4c_5c_6c_7c_1$  has length four; furthermore,  $\varphi(c_4) = \varphi(c_7)$  and  $\varphi(c_5) = \varphi(c_1)$ .*
- (c) *The graph  $G$  contains a path  $c_1uvc_3$  with  $u, v \notin V(C)$ , the cycle  $c_1c_2c_3vu$  bounds a 5-face and every face drawn inside the 8-cycle  $uvc_3c_4c_5c_6c_7c_1$  has length four; furthermore,  $\varphi(c_3) = \varphi(c_6)$ ,  $\varphi(c_2) = \varphi(c_4) = \varphi(c_7)$  and  $\varphi(c_1) = \varphi(c_5)$ .*

By inspection of the cases (a),(b) and (c) of Theorem 16 we obtain the following fact.

**Fact 17.** *Let  $G$  be a plane triangle-free graph with the outer face bounded by a cycle  $C$  of length 7. Suppose that  $G$  is  $C$ -critical and that  $\varphi$  is a precoloring of  $C$  that does not extend to a 3-coloring of  $G$ . Let  $x, y$  and  $z$  be consecutive vertices of  $C$ . If  $\varphi(x) = \varphi(z)$ , then  $y$  is incident with the 5-face of  $G$ .*

We also use the following result dealing with graphs with a triangle.

**Theorem 18** (Aksenov [1]). *Let  $G$  be a plane graph with the outer face bounded by a cycle  $C = c_1c_2\dots$  of length at most 5. If  $G$  is  $C$ -critical and contains exactly one triangle  $T$  distinct from  $C$ , then  $C$  is a 5-cycle, all internal faces of  $G$  other than  $T$  have length exactly four and  $T$  shares at least one edge with  $C$ . Furthermore, if  $C$  and  $T$  share only one edge  $c_1c_2$  and  $\varphi$  is a 3-coloring of  $C$  that does not extend to a 3-coloring of  $G$ , then  $\varphi(c_1) = \varphi(c_3)$  and  $\varphi(c_2) = \varphi(c_5)$ .*

Recall that the notion of a patch was defined in the introduction. First, we need to argue that replacing vertices of degree 3 by critical patches preserves criticality and the number of triangles.

**Lemma 19.** *Let  $G$  be a 4-critical graph, let  $v$  be a vertex of  $G$  of degree 3 and let  $x, y$  and  $z$  be the neighbors of  $v$  in  $G$ . Let  $G'$  be a graph obtained from  $G - v$  by inserting a patch  $P$  with boundary  $C_P = xz'yx'zy'$ , where  $x', y'$  and  $z'$  are new vertices. Then  $G'$  is 4-critical if and only if the patch  $P$  is critical. Furthermore, if  $P$  is critical, then  $t(G') = t(G)$ .*

*Proof.* Clearly, every 4-critical graph has minimum degree at least 3. Furthermore, by Lemma 13 and Theorem 14, if  $\Lambda$  is an open disk bounded by a 4-cycle in a 4-critical plane graph and no triangle is contained in  $\Lambda$ , then  $\Lambda$  is a face. Therefore, if  $G'$  is 4-critical, then  $P$  is a critical patch.

Suppose now conversely that  $P$  is a critical patch. Consider an edge  $e \in E(G')$ . If  $e \notin E(P)$ , then  $e$  is an edge of  $G$  not incident with  $v$ . Since  $G$  is 4-critical, there exists a 3-coloring  $\varphi$  of  $G - e$ . The vertex  $v$  is properly colored, and thus we can by symmetry assume that  $\varphi(x) = \varphi(y)$ . By Theorem 15,  $\varphi$  extends to a 3-coloring of  $P$ , showing that  $G' - e$  is 3-colorable.

On the other hand, consider the case that  $e \in E(P)$ . Since  $G$  is 4-critical, there exists a 3-coloring  $\psi$  of  $G - v$  such that  $\psi(x) = 1$ ,  $\psi(y) = 2$  and  $\psi(z) = 3$ . Suppose that  $\varphi$  does not extend to a 3-coloring of  $P - e$ . If  $e \notin E(C_P)$ , this implies that  $P - e$  contains a  $C_P$ -critical subgraph  $P'$ . By Theorem 14, all faces of  $P'$  distinct from  $C_P$  have length 4. However,  $e$  is

drawn inside one of the faces of  $P'$ , and thus  $P$  would contain a 4-cycle not bounding a face, contrary to the assumption that  $P$  is a critical patch.

Finally, suppose that  $e \in E(C_P)$ , say  $e = xz'$ . Note that all faces of  $P$  have even length, and thus  $P$  is bipartite. Since  $z'$  has degree at least 3 and every 4-cycle in  $P$  bounds a face, we conclude that  $xz'y$  is the only path of length at most two between  $x$  and  $y$  in  $P$ . Let  $P_1$  be obtained from  $P - e$  by adding the edge  $xy$ , and note that  $P_1$  is triangle-free. Let  $C_1 = xyx'zy'$  be the 5-cycle bounding the outer face of  $P_1$ . Since  $\varphi$  does not extend to a 3-coloring of  $P - e$ , it also does not extend to a 3-coloring of  $P_1$ , and thus  $P_1$  contains a  $C_1$ -critical subgraph. This contradicts Theorem 14.

We conclude that  $G' - e$  is 3-colorable for every  $e \in E(G')$ . Since  $G'$  does not contain isolated vertices, this implies that every proper subgraph of  $G'$  is 3-colorable. Suppose that  $G'$  has a proper 3-coloring  $\theta$ . Since  $G$  is not 3-colorable,  $\theta$  cannot be extended to  $v$ , and thus we can assume that  $\theta(x) = 1$ ,  $\theta(y) = 2$  and  $\theta(z) = 3$ . However, that implies that  $\theta(x) = \theta(x')$ ,  $\theta(y) = \theta(y')$  and  $\theta(z) = \theta(z')$ . Since  $\theta$  is a 3-coloring of  $P$ , this contradicts Theorem 15. Therefore,  $G'$  is not 3-colorable, and thus it is 4-critical.

Now we establish a bijection  $f$  between triangles in  $G$  and  $G'$ . If a triangle  $T$  in  $G$  does not contain  $v$ , then  $T$  also appears in  $G'$ , and we set  $f(T) = T$ . If  $T$  contains  $v$ , say  $T = vxy$ , then we set  $f(T) = z'xy$ . Since  $f$  is injective, it suffices to show that it is surjective. Suppose that there exists a triangle  $T' \subset G'$  that is not in the image of  $f$ . Then  $T'$  contains an edge of  $P$ . Since  $P$  is bipartite,  $T'$  contains an edge outside of  $P$ , and since  $x$ ,  $y$  and  $z$  are non-adjacent in the patch  $P$ , we conclude that  $T'$  intersects  $P$  in a path of length two, say  $xwy$ , and  $x$  and  $y$  are adjacent in  $G$ . Since  $f(vxy) = xz'y$  and  $T'$  is not in the image of  $f$ , we conclude that  $w \neq z'$ . Since  $P$  is a critical patch, the 4-cycle  $xz'yw$  bounds a face. However, this implies that  $z'$  has degree two, which is a contradiction. Therefore,  $f$  is indeed a bijection, and thus  $t(G') = t(G)$ .  $\square$

By Lemma 19, the graphs described in Theorem 5 are indeed 4-critical and have exactly 4 triangles. If  $G$  is obtained from a  $\text{Pl}_{4,4f}$ -graph by replacing non-adjacent vertices of degree 3 with (not necessarily critical) patches, then we say that  $G$  is an *expanded  $\text{Pl}_{4,4f}$ -graph*. By Lemma 19, it remains to show that every  $\text{Pl}_4$ -graph is an expanded  $\text{Pl}_{4,4f}$ -graph. First we state several properties of expanded  $\text{Pl}_{4,4f}$ -graphs.

**Claim 20.** *If  $P$  is a patch in an expanded  $\text{Pl}_{4,4f}$ -graph  $G$ , then no vertex of  $P$  is incident with exactly one edge that does not belong to  $P$ .*

*Proof.* Let  $G_0$  be the  $\text{Pl}_{4,4f}$ -graph from which  $G$  is obtained by replacing vertices with patches. Let  $v$  be the vertex of  $G_0$  that is replaced by  $P$  and let  $x, y$  and  $z$  be the neighbors of  $v$ . All vertices of  $V(P) \setminus \{x, y, z\}$  only have neighbors in  $P$ . Each of  $x, y$  and  $z$  has degree at least 3 in  $G_0$ , and thus each of them is incident with at least two edges that are not incident with  $v$ . So, each of  $x, y$  and  $z$  is incident with at least two edges of  $G$  that do not belong to  $P$ .  $\square$

**Claim 21.** *Every face of an expanded  $\text{Pl}_{4,4f}$ -graph  $G$  has length 3, 4 or 5, and every 4-face of  $G$  belongs to a patch.*

*Proof.* Let  $G_0$  be the  $\text{Pl}_{4,4f}$ -graph from which  $G$  is obtained by replacing vertices with patches. By Theorem 4,  $G_0$  belongs to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$ , and thus each face of  $G_0$  has length 3 or 5. Furthermore, observe that replacing a 3-vertex  $v$  by a patch transforms each face of  $G_0$  incident with  $v$  to a face of  $G$  of the same length, whose boundary shares a path of length two with the boundary cycle of  $P$ . Therefore, every face of  $G$  which is not contained in a patch has length 3 or 5.  $\square$

Furthermore, we will use the following simple property of critical graphs.

**Claim 22.** *If  $G$  is a 4-critical graph and  $T$  is a triangle of  $G$ , then  $G - V(T)$  is connected. In particular, if  $G$  is a plane graph, then every triangle in  $G$  bounds a face.*

*Proof.* If  $G - V(T)$  is not connected, then there exist proper subgraphs  $G_1$  and  $G_2$  of  $G$  such that  $G_1 \cup G_2 = G$  and  $G_1 \cap G_2 = T$ . Since  $G$  is 4-critical, both  $G_1$  and  $G_2$  are 3-colorable and by permuting the colors if necessary, we can assume that their 3-colorings match on  $T$ . Together, they would give a 3-coloring of  $G$ , which is a contradiction.  $\square$

A *stretching* of a plane graph  $G$  at a vertex  $w \in V(G)$  is a graph  $G_2$  obtained from  $G$  by the following procedure. Let  $e_1, \dots, e_k$  be the edges incident with  $w$  as drawn in the clockwise order around it. Choose  $m < k$

and let  $G'$  be obtained from  $G$  by removing  $w$ , adding two new vertices  $w_1$  and  $w_2$  and adding edges between  $w_1$  and the endpoints of  $e_1, \dots, e_m$  distinct from  $w$ , and between  $w_2$  and the endpoints of  $e_{m+1}, \dots, e_k$  distinct from  $w$ . Let  $G_1$  be obtained from  $G'$  by either adding a new vertex  $z$  adjacent to  $w_1$  and  $w_2$ , or by adding an edge between  $w_2$  and the endpoint  $z$  of  $e_1$  distinct from  $w$ . Finally, for each face  $f$  of  $G_1$  incident with  $zw_2$ , if  $|f| = 6$ , then replace  $f$  by a quadrangulation, and if  $|f| = 7$ , then replace it by a graph satisfying (a), (b) or (c) of Theorem 16, resulting in the graph  $G_2$ . We call  $G_1$  the *intermediate graph* of the stretching, and the faces of  $G_1$  incident with  $zw_2$  are called *special*.

**Lemma 23.** *Every  $\text{Pl}_4$ -graph is an expanded  $\text{Pl}_{4,4f}$ -graph.*

*Proof.* Assume that  $G$  has the fewest vertices among the  $\text{Pl}_4$ -graphs that are not expanded  $\text{Pl}_{4,4f}$ -graphs. Then  $G$  does not contain any patches, since replacing a patch by a 3-vertex would give a smaller counterexample. We are going to need the following stronger claim.

(1) *Let  $C$  be a 6-cycle in  $G$  and let  $\Delta$  be an open region of the plane bounded by  $C$ , such that all faces in  $\Delta$  have length four. Then  $\Delta$  contains at most one vertex, and if it contains a vertex, then each vertex of  $C$  is incident with an edge that is not drawn in the closure of  $\Delta$ .*

*Proof.* Let  $C = v_1v_2v_3v_4v_5v_6$  and suppose that  $\Delta$  contains at least one vertex, and if it contains only one, then all edges incident with  $v_6$  are drawn in the closure of  $\Delta$ . Let  $G'$  be the graph obtained from  $G$  by removing the vertices in  $\Delta$  (which does not include  $C$ ) and adding a vertex  $v$  adjacent to  $v_1, v_3$  and  $v_5$ . By Theorem 15,  $G'$  is not 3-colorable, since every 3-coloring of  $C$  in which  $v_1, v_3$  and  $v_5$  do not have pairwise distinct colors can be extended to the subgraph of  $G$  drawn in the closure of  $\Delta$ . Therefore,  $G'$  has a 4-critical subgraph  $G''$ . Note that  $|V(G'')| < |V(G)|$ , since if  $\Delta$  contains only one vertex of  $G$ , then  $v_6$  has degree 2 in  $G'$ , and thus  $v_6 \notin V(G'')$ .

Since both  $G$  and  $G''$  are 4-critical,  $G''$  is not a proper subgraph of  $G$ . Hence  $v$  and all edges incident to it belong to  $G''$ . Triangles in  $G''$  that are not in  $G$  can only be created by adding  $v$  to an edge, say  $v_1v_3$ ; in this case,  $v_1v_2v_3$  is a triangle in  $G$  which bounds a face (by Claim 22), and thus  $v_2$  has degree two in  $G'$  and does not belong to  $G''$ , which means that creating a new triangle in  $G''$  destroys another triangle. Thus  $t(G'') \leq t(G) = 4$ . Since  $G''$  is not 3-colorable, it contains exactly four triangles. By the minimality

of  $G$ , the graph  $G''$  is an expanded  $\text{Pl}_{4,4f}$ -graph and by Claim 21, each face of  $G''$  has length at most 5. Let  $G_1$  be the graph obtained from  $G'' - v$  by adding the subgraph  $H$  of  $G$  contained in the closure of  $\Delta$ . Then  $G_1$  is a subgraph of  $G$ .

If at least two faces of  $G''$  incident with  $v$  have length 4, then  $v$  is a vertex of a patch  $P''$  of  $G''$  and all neighbors of  $v$  belong to  $P''$ . We claim that  $P = (P'' - v) + H$  is a patch; this is clear if  $v$  is incident with three 4-faces in  $G''$ . If  $v$  is incident with exactly two 4-faces, then let  $f$  be the face incident with  $v$  of length other than four. Note that at least one edge  $e$  of  $C$  does not belong to  $G''$  and in  $G$ , it is drawn in the region of the plane corresponding to  $f$ . By symmetry, we can assume that  $e = v_1v_6$ . Since all faces of  $G''$  have length at most 5, if  $v_6 \in V(G'')$ , then  $v_1$  and  $v_6$  would be joined by a path  $Q$  of length two in  $G''$ . The path  $Q$  together with the edge  $v_1v_6$  would form a triangle in  $G$  which does not correspond to any triangle in  $G''$ , and thus  $G''$  would only have at most three triangles, which is a contradiction. Therefore,  $v_6 \notin V(G'')$ , and thus all neighbors of  $v_6$  in  $G_1$  belong to  $P$ , showing that  $P$  is a patch. Consequently,  $G_1$  is an expanded  $\text{Pl}_{4,4f}$ -graph.

Hence, we can assume that at most one face of  $G''$  incident with  $v$  has length four. If exactly one face incident with  $v$  had length 4, then  $v$  would belong to a patch  $P$  and it would be incident with exactly one edge not belonging to  $P$ , contradicting Claim 20. Therefore, no face incident with  $v$  has length four, and as in the previous paragraph, we conclude that  $H$  is a patch in  $G_1$  that replaces the vertex  $v$  of degree 3 of  $G''$ . Again, it follows that  $G_1$  is an expanded  $\text{Pl}_{4,4f}$ -graph.

Since  $G_1$  is not 3-colorable and  $G$  is 4-critical,  $G = G_1$  and  $G$  is an expanded  $\text{Pl}_{4,4f}$ -graph. This is a contradiction.  $\square$

**(2)**  $G$  does not contain separating 4-cycles.

*Proof.* Suppose that  $C = v_1v_2v_3v_4$  is a separating 4-cycle in  $G$ . Let  $\Lambda_1$  and  $\Lambda_2$  be the two connected regions obtained from the plane by removing  $C$ . For  $i \in \{1, 2\}$ , let  $G_i$  be the subgraph of  $G$  drawn in the closure of  $\Lambda_i$ . By Lemma 13,  $G_i$  is  $C$ -critical, and by Theorem 18,  $G_i$  contains at least two triangles. Since  $t(G) = 4$ , we conclude that  $t(G_1) = t(G_2) = 2$ .

For  $i, j \in \{1, 2\}$ , let  $G_{i,j}$  be the graph obtained from  $G_i$  by adding the edge  $v_jv_{j+2}$ . Note that  $C + v_jv_{j+2}$  has a unique 3-coloring up to permutation of colors. Therefore, any 3-colorings of  $G_{1,j}$  and  $G_{2,j}$  could be combined to



a 3-coloring of  $G$ . We conclude that at least one of  $G_{1,j}$  and  $G_{2,j}$  is not 3-colorable. By symmetry, we can assume that  $G_{1,1}$  is not 3-colorable.

Let  $G'_{1,1}$  be a 4-critical subgraph of  $G_{1,1}$ . By Claim 22, each triangle in  $G'_{1,1}$  bounds a face, and thus  $v_1v_3$  belongs to at most two triangles of  $G'_{1,1}$ . Since  $t(G_1) = 2$ , it follows that  $t(G'_{1,1}) \leq 4$ . Since  $G'_{1,1}$  is 4-critical, it follows that  $t(G'_{1,1}) = 4$ , i.e.,  $G'_{1,1}$  is a  $\text{Pl}_{4,f}$ -graph, and that  $v_1v_3$  belongs to two triangles of  $G'_{1,1}$ . Let  $v_1v'_2v_3$  and  $v_1v'_4v_3$  be the triangles incident with  $v_1v_3$  labelled so that for  $k \in \{2, 4\}$ ,  $v_k$  is either equal to  $v'_k$  or it is drawn in  $G_{1,1}$  in the region of the plane corresponding to the face of  $G'_{1,1}$  bounded by  $v_1v'_kv_2$ .

Since  $C$  is a separating cycle,  $|V(G'_{1,1})| < |V(G)|$ , and by the minimality of  $G$ , it follows that  $G'_{1,1}$  is an expanded  $\text{Pl}_{4,4f}$ -graph. By Claim 21, every face of  $G'_{1,1}$  has length at most five. Consider a face  $f$  of  $G'_{1,1}$  not incident with  $v_1v_3$ . Since the subgraph of  $G$  drawn in the closure of  $f$  contains no triangles, by Lemma 13 and Theorem 14,  $f$  is a face of  $G$  as well. Furthermore, if  $v_k \neq v'_k$  for some  $k \in \{2, 4\}$ , then Lemma 13 and Theorem 14 imply that  $v_1v'_kv_3v_k$  is a face of  $G$  and  $v_k$  has degree two in  $G_{1,1}$ . It follows that  $V(G_{1,1}) \setminus V(G'_{1,1}) \subseteq \{v_2, v_4\}$  and each vertex  $v \in V(G_{1,1}) \setminus V(G'_{1,1})$  is adjacent in  $G_{1,1}$  only to  $v_1$  and  $v_3$ .

Suppose that  $G'_{1,1}$  has a 4-face. In that case, it contains a patch  $P$ . Let  $x, y$  and  $z$  be the vertices of the boundary cycle  $K$  of  $P$  that have neighbors only in  $P$ . Note that  $v_1v_3 \notin E(K)$ , since both faces incident with  $v_1v_3$  are triangles. Furthermore, at most two of  $x, y$  and  $z$  are incident with  $C$ , since they form an independent set. Therefore, we can assume that all edges incident with  $x$  in  $G$  belong to  $P$ . This contradicts (1). We conclude that  $G'_{1,1}$  has no 4-faces, and thus it belongs to  $\mathcal{TW} \cup \mathcal{TW}_1 \cup \mathcal{TW}_2$  by Theorem 4.

Consequently, both vertices of the diamond edge  $v_1v_3$  of  $G'_{1,1}$  have degree exactly 3 in  $G'_{1,1}$ . Since  $G'_{1,1}$  is 4-critical, there exists a 3-coloring  $\varphi$  of  $G'_{1,1} - \{v_1, v_3\}$  and  $\varphi(v'_2) \neq \varphi(v'_4)$ . Set  $\varphi(v_k) = \varphi(v'_k)$  for  $k \in \{2, 4\}$  and note that  $\varphi$  is a proper 3-coloring of  $G_{1,2}$ . Since  $G$  is not 3-colorable, it follows that  $G_{2,2}$  is not 3-colorable. By an argument symmetrical to the one for  $G_{1,1}$ , we conclude that  $G_{2,2}$  contains a  $\text{Pl}_{4,4f}$ -graph  $G'_{2,2}$  as a subgraph such that  $V(G_{2,2}) \setminus V(G'_{2,2}) \subseteq \{v_1, v_3\}$  and each vertex  $v \in V(G_{2,2}) \setminus V(G'_{2,2})$  is only adjacent to  $v_2$  and  $v_4$  in  $G_{2,2}$ .

Suppose that  $G'_{1,1} \neq G_{1,1}$ , and thus say  $v_2 \in V(G_{1,1}) \setminus V(G'_{1,1})$  is adjacent in  $G_1$  only to  $v_1$  and  $v_3$ . Since  $v_2v_4$  is a diamond edge of  $G'_{2,2}$ , vertex  $v_2$  has degree 3 in  $G'_{2,2}$ . If  $G'_{2,2} = G_{2,2}$ , this would imply that  $v_1$  and  $v_3$  are the only neighbors of  $v_2$  in  $G_2$ , and thus  $v_2$  would have degree two in  $G$ ,

contrary to the assumption that  $G$  is 4-critical. Hence, we can assume that say  $v_1 \in V(G_{2,2}) \setminus V(G'_{2,2})$  is adjacent in  $G_2$  only to  $v_2$  and  $v_4$ . Since  $G'_{1,1}$  and  $G'_{2,2}$  are 4-critical, there exist 3-colorings  $\varphi_1$  of  $G'_{1,1} - v_1v_3$  and  $\varphi_2$  of  $G'_{2,2} - v_2v_4$  such that  $\varphi_1(v_1) = \varphi_1(v_3) = 1$ ,  $\varphi_1(v'_2) = 2$ ,  $\varphi_1(v'_4) = 3$ ,  $\varphi_2(v_2) = \varphi_2(v_4) = 3$ ,  $\varphi_2(v'_1) = 2$  and  $\varphi_2(v'_3) = 1$ . Then each vertex of  $G$  is colored by  $\varphi_1$  or  $\varphi_2$  and if a vertex ( $v_3$  or  $v_4$ ) belongs to both  $G'_{1,1}$  and  $G'_{2,2}$ , then it is assigned the same color by  $\varphi_1$  and  $\varphi_2$ . Thus the union of  $\varphi_1$  and  $\varphi_2$  gives a 3-coloring of  $G$ , which is a contradiction.

Therefore,  $G'_{1,1} = G_{1,1}$ , and by symmetry,  $G'_{2,2} = G_{2,2}$ . It follows that  $G$  has no 4-face, and thus  $G$  is a  $\text{Pl}_{4,4f}$ -graph. This is a contradiction.  $\square$

**(3)** *If  $K$  is a separating 5-cycle in  $G$  and  $\Delta$  is a region of the plane bounded by  $K$  such that  $\Delta$  contains at most one triangle, then  $\Delta$  contains exactly one vertex, which has three neighbors in  $K$ .*

*Proof.* By Theorem 14 and the criticality of  $G$ ,  $\Delta$  contains a triangle  $T$ , as  $\Delta$  is not a face of  $G$ . Since  $K$  is separating, Theorem 18 implies that  $T$  shares exactly one edge with  $K$ . Let  $\{z\} = V(T) \setminus V(K)$ . Note that  $T \cup K$  contains a 6-cycle, bounding an open region  $\Delta_0 \subset \Delta$ . Since all edges incident with  $z$  are drawn in the closure of  $\Delta_0$ , (1) implies that  $\Delta_0$  contains no vertices, and thus  $z$  is the only vertex in  $\Delta$ .  $\square$

**(4)** *The following configuration does not appear in  $G$ : a path  $z_1z_2z_3$  and a vertex  $z$  of degree 3 adjacent to  $z_1$ ,  $z_2$  and  $z_3$ .*

*Proof.* Since  $G$  does not contain separating triangles, if  $z_1$  is adjacent to  $z_3$ , then  $G = K_4$ . This is a contradiction, since  $G$  is not a  $\text{Pl}_{4,4f}$ -graph.

Let  $G'$  be the graph obtained from  $G - z$  by identifying  $z_1$  with  $z_3$  to a new vertex  $u$ . Since  $G$  is not 3-colorable,  $G'$  also is not 3-colorable. Let  $G''$  be a 4-critical subgraph of  $G'$ . Note that  $G''$  has at least four triangles, and since the triangles  $zz_1z_2$  and  $zz_2z_3$  disappear during the construction of  $G'$ , we conclude that  $G''$  contains at least two triangles  $ux_1x_2$  and  $uy_1y_2$  such that  $z_1x_1x_2z_3$  and  $z_1y_1y_2z_3$  are (not necessarily disjoint) paths between  $z_1$  and  $z_3$  in  $G - \{z, z_2\}$ . Since  $G''$  contains no separating triangles by Claim 22, we conclude that  $G''$  contains exactly two such triangles, and thus  $t(G'') = 4$ . Let  $\Delta_1$  be the open disk bounded by  $z_1z_3z_2x_1$  in  $G$  corresponding to the

face  $ux_1x_2$  of  $G'$ , and let  $\Delta_2$  be the open disk bounded by  $z_1z_2z_3y_2y_1$  in  $G$  corresponding to the face  $uy_1y_2$  of  $G'$ . By swapping the labels of  $x_i$  and  $y_i$  (for  $i \in \{1, 2\}$ ) if necessary, we can assume that  $\Delta_1$  and  $\Delta_2$  are disjoint. Note that since  $t(G'') = 4$ , neither  $\Delta_1$  nor  $\Delta_2$  contains a triangle of  $G$ , and thus  $\Delta_1$  and  $\Delta_2$  are faces of  $G$  by (3).

By the minimality of  $G$ , it follows that  $G''$  is an expanded  $\text{Pl}_{4,4f}$ -graph, and all faces of  $G''$  have length at most 5. By Lemma 13 and Theorem 14, all faces of  $G''$  other than  $ux_1x_2$  and  $uy_1y_2$  are also faces of  $G$ . In particular, since  $G$  does not contain patches,  $G''$  also does not contain patches, and thus  $G''$  has no 4-faces. It follows that  $G$  has no 4-faces. Therefore,  $G$  is a  $\text{Pl}_{4,4f}$ -graph, which is a contradiction.  $\square$

(5) *The following configuration does not appear in  $G$ : a triangle  $T = z_1z_2z_3$  such that all vertices of  $T$  have degree 3 and  $z_3$  is adjacent to a vertex  $x_3$  distinct from  $z_1$  and  $z_2$  that has degree 3 and belongs to a triangle.*

*Proof.* Let  $x_1$  and  $x_2$  be the neighbors of  $z_1$  and  $z_2$ , respectively, outside of  $T$ . By (4), we have  $x_1 \neq x_2 \neq x_3 \neq x_1$ . Furthermore, the vertices  $x_1, x_2$  and  $x_3$  form an independent set in  $G$ , as otherwise every 3-coloring of  $G - V(T)$  would extend to  $G$ , contrary to the 4-criticality of  $G$ . Let  $G'$  be the graph obtained from  $G - V(T) - x_3$  by adding the edge  $x_1x_2$ . Note that every 3-coloring of  $G'$  extends to a 3-coloring of  $G$ , and thus  $G'$  is not 3-colorable. Let  $G''$  be a 4-critical subgraph of  $G$ . Note that  $G''$  contains at least four triangles, and since  $T$  as well as the triangle incident with  $x_3$  disappear during the construction of  $G'$ , it follows that  $x_1x_2$  belongs to at least two triangles in  $G''$ . By Claim 22,  $x_1x_2$  belongs to exactly two triangles, each of them bounding a face of  $G''$ . Therefore,  $G$  contains a 4-cycle  $x_1ux_2v$  separating two of its triangles from  $T$ . By (2), it follows that  $u$  is adjacent to  $v$ . By (3), either  $x_1z_1z_2x_2u$  or  $x_1z_1z_2x_2v$  bounds a face, and thus  $u$  or  $v$  has degree 3. This contradicts (4).  $\square$

Let  $v_1v_2v_3v_4$  be a 4-face in  $G$ , if possible chosen so that it contains two adjacent vertices that are only incident with 4-faces. Since  $G$  contains no separating triangles,  $v_1v_3$  and  $v_2v_4$  are not edges. Suppose that  $v_1$  and  $v_3$  are joined by a path  $v_1x_1x_2v_3 \subset G - \{v_2, v_4\}$  of length 3, and that  $v_2$  and  $v_4$  are joined by a path  $v_2y_1y_2v_4 \subset G - \{v_1, v_3\}$  of length 3. By symmetry and

planarity, we can assume that  $x_1 = y_1$ . If both  $v_1$  and  $v_2$  have degree 3, then every 3-coloring of  $G - \{v_1, v_2\}$  extends to  $G$ , contrary to the assumption that  $G$  is 4-critical. By symmetry,  $v_1$  has degree at least 4, and since  $v_4y_2y_1v_1$  is not a separating 4-cycle,  $v_1$  is adjacent to  $y_2$ . By (3) applied to the 5-cycle  $v_4v_1v_2y_1y_2$ , we have  $x_2 = y_2$ . But then  $V(G) = \{v_1, v_2, v_3, v_4, x_1, x_2\}$  and  $G$  is 3-colorable.

Therefore, we can by symmetry assume that  $v_1$  and  $v_3$  are not joined by a path of length 3 in  $G - \{v_2, v_4\}$ . Let  $G'$  be the graph obtained from  $G$  by identifying  $v_1$  and  $v_3$  to a single vertex  $w$ . Clearly,  $G'$  has exactly the four triangles that originally belonged to  $G$  (possibly with  $v_1$  or  $v_3$  relabelled to  $w$ ). Since every 3-coloring of  $G'$  gives a 3-coloring of  $G$ , we conclude that  $G'$  is not 3-colorable. Let  $G''$  be a 4-critical subgraph of  $G'$ . By the minimality of  $G$ ,  $G''$  is an expanded  $\text{Pl}_{4,4f}$ -graph. In particular, all faces of  $G''$  have length at most 5. For a face  $f$  of  $G''$ , let  $C_f$  denote the corresponding cycle in  $G$  (either equal to  $f$  up to relabelling of  $w$  to  $v_1$  or  $v_3$ , or obtained from  $f$  by replacing  $w$  by the path  $v_1v_2v_3$ ). Since all triangles of  $G$  are faces of  $G''$ , Theorem 14 implies that if  $|C_f| = |f|$ , then  $C_f$  is a face of  $G$ . Furthermore, if  $|C_f| = |f| + 2$ , then  $4 \leq |f| \leq 5$  and  $C_f$  does not bound a face of  $G$ , since  $v_2$  has degree at least 3. Let  $G_f$  denote the subgraph of  $G$  drawn in the region of the plane bounded by  $C_f$  and corresponding to  $f$ , and note that  $G_f$  is one of the graphs described by Theorem 14 or Theorem 16. Therefore, the following holds.

**(6)** *The graph  $G$  is obtained by stretching from an expanded  $\text{Pl}_{4,4f}$ -graph  $G_0$  at a vertex  $w_0$ , where  $G$  and  $G_0$  have the same triangles.*

In particular, all faces of  $G$  have length at most 5. By the choice of the 4-face  $v_1v_2v_3v_4$ , we can also assume that the following property is satisfied.

**(7)** *Let  $G_0$  and  $w_0$  be as in (6). Let  $v_1v_2v_3$  be the path replacing  $w_0$  in the intermediate graph of the stretching. The path  $v_1v_2v_3$  is contained in the boundary of a 4-face in  $G$ . Furthermore, if  $G$  contains an edge whose vertices are only incident with 4-faces, then  $v_1$  is only incident with 4-faces.*

Let  $G_0$  and  $w_0$  satisfying (6) be chosen so that if  $G_0$  contains at least one patch, then (7) is satisfied, and subject to that with  $|V(G_0)|$  as small as possible. Observe that  $G_0$  does not necessarily have to be created from  $G$  by contracting a 4-face. Let  $G_1$  be the intermediate graph of the stretching, let  $f_1$  and  $h_1$  be its special faces and let  $f_0$  and  $h_0$  be the corresponding faces of

$G_0$ .

(8) *The graph  $G_0$  has no 4-faces.*

*Proof.* Suppose that  $G_0$  has a 4-face, and thus it contains a patch  $P_0$  bounded by a 6-cycle  $C_0 = z_1z_2z_3z_4z_5z_6$ , where  $z_2, z_4$  and  $z_6$  only have neighbors in  $P_0$ . Let  $r_2, r_4$  and  $r_6$  be the faces of  $G_0$  incident with  $z_2, z_4$  and  $z_6$ , respectively, whose length is not four. Since  $G_0$  is 2-connected, these faces are pairwise distinct. If  $C_0$  is a 6-cycle in  $G$ , then the open region  $\Delta$  corresponding to  $P_0$  contains only 4-faces and at least one vertex of  $G$  lies inside  $\Delta$ . There exists  $k \in \{2, 4, 6\}$  such that  $f_0 \neq r_k \neq h_0$ , and thus all edges incident with  $z_k$  in  $G$  are drawn in the closure of  $\Delta$ . This contradicts (1); hence,  $C_0$  contains  $w_0$  and corresponds to an 8-cycle  $C_1$  in  $G_1$  containing the path  $v_1v_2v_3$ . Note that all faces of  $G$  drawn in the open region  $\Lambda$  of the plane bounded by  $C_1$  and corresponding to  $P_0$  have length 4. By symmetry, we can assume that  $w_0 \in \{z_1, z_2\}$  and  $f_0$  is contained in the patch  $P_0$ .

Suppose that  $h_0$  does not share an edge with  $C_0$ ; then,  $w_0 = z_1, r_2$  and  $r_6$  are faces of  $G$  and  $v_1$  is incident with one of them. Since neither  $r_2$  nor  $r_6$  is a 4-face, (7) implies that  $G$  does not contain an edge whose vertices are only incident with 4-faces. Hence,  $|V(P_0) \setminus V(C_0)| = 1$  and the only neighbors of  $z_1$  in  $P_0$  are  $z_2$  and  $z_6$ . By (7),  $G$  contains a 4-face  $v_1v_2v_3z$ . If  $z$  lies inside  $\Lambda$ , then note that it is adjacent neither to  $z_2$  nor to  $z_6$ , and thus  $z$  has a neighbor inside  $\Lambda$ . If  $z$  lies outside  $\Lambda$ , then similarly  $v_2$  has a neighbor inside  $\Lambda$ . In both cases  $G$  contains an edge whose vertices are incident only with 4-faces, a contradiction. Therefore,  $h_0$  shares an edge with  $C_0$  (we can assume that  $h_0 = r_2$ ). Since the same conclusions would have to hold for any other patch in  $G_0$ , the only patch in  $G_0$  is  $P_0$ .

Label the vertices of  $C_1$  so that the vertices of  $V(C_0) \setminus \{w_0\}$  retain their labels and  $C_1 = z_1xyz_2z_3z_4z_5z_6$ . Note that all edges incident with  $y, z_2, z_4$  and  $z_6$  in  $G_1$  are drawn in the closure of  $\Lambda$ . Also, either  $x$  is adjacent to  $z_1$  in  $G_0$  and  $|h_1| = |h_0| = 5$ , or  $|h_0| = 5$  and  $|h_1| = 7$  and all edges incident with  $x$  are drawn in the closure of  $\Lambda$ . Let  $h_0 = z_1z_2z_3ab$ , where  $b$  may be equal to  $x$ . Since  $C = z_1baz_3z_4z_5z_6$  is a 7-cycle in  $G$  whose interior (the part of the plane bounded by  $C$  and containing  $\Lambda$ ) does not contain any triangle, it satisfies (b) or (c) of Theorem 16. Let  $G'_0$  be the graph obtained from  $G_0$  by replacing the patch  $P_0$  with a 3-vertex  $z'$  adjacent to  $z_1, z_3$  and  $z_5$ . Since  $G_0$  has only one patch,  $G'_0$  has no patches and it is a  $\text{Pl}_{4,4f}$ -graph. Furthermore,  $G$  is obtained from  $G'_0$  by stretching at vertex  $z'$  (split out the edge  $z'z_3$ , then




Figure 6: Possibilities for a fragment  $F$ .

make its new endvertex adjacent to  $z_5$  and relabel the vertices created from  $z'$  by  $z_4$  and  $z_6$ ). Since  $|V(G'_0)| < |V(G_0)|$ , this is a contradiction with the choice of  $G_0$ .  $\square$

Suppose that  $\{a, c\}$  is a vertex-cut in  $G_0$  and  $F$  is an induced subgraph of  $G_0$  such that all edges of  $G_0$  incident with  $V(F) \setminus \{a, c\}$  belong to  $F$ . We say that  $F$  is a *fragment with attachments  $a$  and  $c$*  if  $F$  either isomorphic to the Havel's quasiedge  $H_0$  and  $a$  and  $c$  are its vertices of degree two, or if  $F$  consists of a 4-cycle  $au_1u_2u_3$  and edges  $u_1u_3$  and  $u_2c$ . See Figure 6 for an illustration. Since all triangles of  $G_0$  belong to  $G$  and stretching does not preserve size of at least one face,  $G_0$  is not  $K_4$ , and thus  $G_0$  contains two fragments.

Since stretching has at most two special faces and fragment contains configurations forbidden by (4) and (5), the stretching must occur at a vertex of each of the fragments. Hence,  $G_0$  does not have two vertex-disjoint fragments, and we conclude that  $G$  is one of the graphs  $M$  and  $K'_4$  depicted in Figures 2 and 3. The possible intermediate graphs  $G_1$  are drawn in Figure 7. Thus we have the three cases below.

Case 1:  $G_1 = X_1$ . Let  $\psi$  be a 3-coloring of  $G_1$  such that  $\psi(v_1) = \psi(x) = \psi(v_4) = 1$ ,  $\psi(v_2) = \psi(x_1) = \psi(x_2) = 2$  and  $\psi(v_3) = \psi(x_3) = \psi(x_4) = 3$ . This coloring does not extend to a 3-coloring of  $G$ , and by symmetry, we can assume that it does not extend to a 3-coloring of  $G_{h_0}$ . By Fact 17, it follows that  $x$ ,  $x_3$  and  $v_4$  are incident with a common 5-face in  $G$ , and thus  $x_3$  has degree 3. This contradicts (4).

Case 2:  $G_1 = X_2$ . Let  $\psi$  be a 3-coloring of  $G_1$  such that  $\psi(v_1) = \psi(x) = \psi(y_1) = 1$ ,  $\psi(v_2) = \psi(x_1) = \psi(y_2) = 2$  and  $\psi(v_3) = \psi(x_2) = \psi(y) = 3$ . By symmetry,  $\psi$  does not extend to  $G_{h_0}$ , and by Theorem 16,  $x_2$  is adjacent to  $v_3$  and  $xx_2v_3y_2y$  is a 5-face. However, then  $y_2$  has degree 3 and we again obtain a contradiction with (4).

Case 3:  $G_1 = Y_1$ . Let  $\psi$  be a 3-coloring of  $G_1$  such that  $\psi(v_1) = \psi(x_1) =$




Figure 7: Stretchable  $\text{Pl}_{4,4f}$ -graphs.

$\psi(y_4) = 1$ ,  $\psi(v_2) = \psi(y_2) = \psi(x_2) = \psi(y_3) = 2$  and  $\psi(v_3) = \psi(x) = \psi(y_1) = 3$ . If  $\psi$  does not extend to a 3-coloring of  $G_{f_0}$ , then by Fact 17,  $G$  contains a 5-face incident with  $v_3$ ,  $x_2$  and  $x$ ; hence,  $x_2$  has degree 3 and contradicts (4). Since  $G$  is not 3-colorable, it follows that  $\psi$  does not extend to a 3-coloring of  $G_{h_0}$ . By Fact 17,  $G$  contains a 5-face incident with  $v_1$ ,  $y_3$  and  $x$ , and by Theorem 14,  $y_4$  is incident with the 5-face as well. However, then both  $y_3$  and  $y_4$  have degree 3 in  $G$ , which contradicts (5).

This finishes the proof of Lemma 23.  $\square$

*Proof of Theorem 5.* By Lemma 19, every graph obtained from a  $\text{Pl}_{4,4f}$ -graph by replacing non-adjacent 3-vertices with critical patches is 4-critical and has exactly four triangles. Conversely, if  $G$  is a  $\text{Pl}_4$ -graph, then it is an expanded  $\text{Pl}_{4,4f}$ -graph by Lemma 23, and since  $G$  is 4-critical, all the patches in  $G$  are critical by Lemma 19.  $\square$

## References

- [1] AKSENOV, V. A. The extension of a 3-coloring on planar graphs. *Diskret. Analiz*, Vyp. 26 Grafy i Testy (1974), 3–19, 84.
- [2] AKSENOV, V. A. Private communication, 1976.
- [3] AKSENOV, V. A., BORODIN, O. V., AND GLEBOV, A. N. Continuation of a 3-coloring from a 6-face onto a plane graph without 3-cycles. *Diskretn. Anal. Issled. Oper. Ser. 1 10* (2003), 3–11. in Russian.
- [4] AKSENOV, V. A., BORODIN, O. V., AND GLEBOV, A. N. Continuation of a 3-coloring from a 7-face onto a plane graph without 3-cycles. *Sib. Elektron. Mat. Izv. 1* (2004), 117–128. in Russian.
- [5] AKSIONOV, V., AND MEL'NIKOV, L. Essay on the theme: The three-color problem. *Combinatorics, Keszthely 1976*, Colloq. Math. Soc. Janos Bolyai 18, 23–34 (1978), 1978.
- [6] AKSIONOV, V. A., AND MELNIKOV, L. S. Some counterexamples associated with the three-color problem. *Combin. Theory Ser. B 28* (1980), 1–9.
- [7] BORODIN, O. V. A new proof of Grünbaum's 3 color theorem. *Discrete Math. 169* (1997), 177–183.
- [8] BORODIN, O. V. Colorings of plane graphs: A survey. *Discrete Math. 313* (2013), 517–533.
- [9] DVOŘÁK, Z., KRÁL', D., AND THOMAS, R. Coloring planar graphs with triangles far apart. submitted, 2009.
- [10] DVOŘÁK, Z., KRÁL', D., AND THOMAS, R. Coloring triangle-free graphs on surfaces. In *Proceedings of the twentieth Annual ACM-SIAM Symposium on Discrete Algorithms* (Philadelphia, PA, USA, 2009), SODA '09, Society for Industrial and Applied Mathematics, pp. 120–129.



- [11] DVOŘÁK, Z., KRÁL', D., AND THOMAS, R. Three-coloring triangle-free graphs on surfaces I. Extending a coloring to a disk with one triangle. *ArXiv e-prints 1010.2472* (Oct. 2010).
- [12] DVOŘÁK, Z., AND LIDICKÝ, B. 3-coloring triangle-free planar graphs with a precolored 8-cycle. *ArXiv e-prints 1305.2467* (May 2013).
- [13] GIMBEL, J., AND THOMASSEN, C. Coloring graphs with fixed genus and girth. *Transactions of AMS 349* (1997), 4555–4564.
- [14] GRÖTZSCH, H. Ein Dreifarbenatz für Dreikreisfreie Netze auf der Kugel. *Math.-Natur. Reihe 8* (1959), 109–120.
- [15] GRÜNBAUM, B. Grötzsch's theorem on 3-coloring. *Michigan Math. J. 10* (1963), 303–310.
- [16] HAVEL, I. On a conjecture of Grünbaum. *Journal of Combinatorial Theory, Series B 7* (1969), 184–186.
- [17] KOSTOCHKA, A. V., AND YANCEY, M. A Brooks-type result for sparse critical graphs. submitted, 2012.
- [18] SACHS, H. Einführung in die theorie der endlichen graphen. teil 2. Mathematisch-Naturwissenschaftliche Bibliothek 44. Leipzig: BSB B. G. Teubner Verlagsgesellschaft. 272 S. m. 151 Abb., M 25.00 (1972), 1972.
- [19] STEINBERG, R. The state of the three color problem. Quo Vadis, Graph Theory? *Ann. Discrete Math. 55* (1993), 211–248.
- [20] THOMAS, R., AND WALLS, B. Three-coloring Klein bottle graphs of girth five. *J. Combin. Theory Ser. B 92* (2004), 115–135.